

ELLIPTIC SYSTEMS WITH BOUNDED SOLUTIONS AND DOUBLE PHASE FUNCTIONALS

SISTEMI ELLITTICI CON SOLUZIONI LIMITATE E FUNZIONALI DOPPIA FASE

FRANCESCO LEONETTI

ABSTRACT. We study boundedness of weak solutions to elliptic systems of partial differential equations in divergence form, under the so-called p, q growth conditions. Examples are obtained by writing the Euler system of some double phase functionals.

SUNTO. Studiamo la limitatezza delle soluzioni deboli di sistemi di equazioni differenziali di tipo ellittico in forma di divergenza, sotto le cosiddette condizioni di crescita p, q . Esempi sono ottenuti scrivendo l'equazione di Eulero di alcuni funzionali doppia fase.

2010 MSC. Primary 35J47; Secondary 35B65.

KEYWORDS. Elliptic, system, bounded, weak, solution, double, phase, p, q growth.

1. ELLIPTIC SYSTEMS WITH BOUNDED SOLUTIONS

We consider the system of partial differential equations in divergence form

$$(1) \quad \left\{ \begin{array}{l} - \sum_{i=1}^n D_i [A_i^1(x, Du(x))] = 0, \\ - \sum_{i=1}^n D_i [A_i^2(x, Du(x))] = 0, \\ \dots\dots\dots \\ - \sum_{i=1}^n D_i [A_i^m(x, Du(x))] = 0, \end{array} \right.$$

Bruno Pini Mathematical Analysis Seminar, Vol. 15 No. 1 (2024) pp. 138-163
Dipartimento di Matematica, Università di Bologna
ISSN 2240-2829.

where $x \in \Omega \subset \mathbb{R}^n$, Ω is bounded and open, $u : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}^m$ and u^1, u^2, \dots, u^m are the components of u , so that, $u = (u^1, u^2, \dots, u^m)$. We assume that u is a weak solution of the previous system, $A(x, z)$ is measurable with respect to x and continuous with respect to z . We ask what are sufficient conditions on $A(x, z)$ that force u to be bounded. We first consider the scalar case $m = 1$, that is, the case in which the system consists of only one equation

$$(2) \quad - \sum_{i=1}^n D_i [A_i^1(x, Du(x))] = 0,$$

where $u : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}$ and, for some constants $p \in (1, +\infty)$, $c_1, \nu \in (0, +\infty)$, $c_2 \in [0, +\infty)$,

$$(3) \quad |A_i^1(x, z)| \leq c_1(|z|^{p-1} + 1),$$

$$(4) \quad \nu|z|^p - c_2 \leq \sum_{i=1}^n A_i^1(x, z)z_i.$$

Let us assume that $u \in W^{1,p}(\Omega; \mathbb{R})$ is a weak solution of equation (2); then

$$u \in C_{loc}^{0,\alpha}(\Omega; \mathbb{R}).$$

This is the celebrated regularity result obtained by De Giorgi, Nash, Moser, at the end of '50, see [55], [108], [107].

In the vectorial case $m \geq 2$, singular solutions to elliptic systems may appear, as De Giorgi (1968) shows in [56]; see also [101], [75], [67], [110], [72], [85], [113], [74], [84], [88], [78], [77], [115], [68], [105]. See also the reviews [102], [103], [87].

Let us come back to the scalar case $m = 1$ and let us note that the exponent p of the growth condition from above (3) is the same exponent in the growth condition from below (4) and it is the same exponent of the Sobolev space $W^{1,p}(\Omega)$ to which the weak solution

u is assumed to belong. If we allow a bigger exponent q in the growth condition from above, namely

$$(5) \quad |A_i^1(x, z)| \leq c_1(|z|^{q-1} + 1),$$

$$(6) \quad \nu|z|^p - c_2 \leq \sum_{i=1}^n A_i^1(x, z)z_i,$$

$$p < q,$$

then, singular solutions may appear: Giaquinta, Marcellini, Hong, end of '80, see [73], [96], [82].

Some regularity can be obtained if q is close to p : [97], [98], [106], [69], [116], [14], [70], [94], [29], [30], [1], [114], [82], [11], [28], [43] [62], [63], [71], [25], [12], [92], [32], [38], [15], [26], [7], [36], [16], [23], [34], [57], [27], [39], [60], [21], [2], [3], [8], [9], [81], [46], [22], [20], [109], [17], [5], [99], [58], [59], [100], [48], [49], [52], [83] [54], [40]. See also the reviews [90], [102], [104],

Let us come back to the vectorial case $m \geq 1$, that is, we consider the system (1), where $u : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}^m$ and, for constants $q, p \in (1, +\infty)$, $c_1, \nu \in (0, +\infty)$, $c_2 \in [0, +\infty)$,

$$(7) \quad |A_i^\alpha(x, z)| \leq c_1(|z|^{q-1} + 1),$$

$$(8) \quad \nu|z^\alpha|^p - c_2 \leq \sum_{i=1}^n A_i^\alpha(x, z)z_i^\alpha,$$

$$(9) \quad 1 < p < q < \frac{p^*}{p} \frac{n}{n+1}, \quad p < n.$$

Assume that

$$(10) \quad u \in W^{1,q}(\Omega; \mathbb{R}^m)$$

is a weak solution to system (1), then

$$(11) \quad u \in L_{loc}^{\infty}(\Omega; \mathbb{R}^m),$$

see Cupini-Leonetti-Mascolo (2022): [37].

In order to get bounded solutions, we need to keep away De Giorgi's counterexample, [56]. So, we must assume some structure condition that excludes De Giorgi's system. In the previous result, such a structure condition is (8): let us call it "componentwise coercivity". It says that the α row of the system $A_i^{\alpha}(x, Du)$ contains the gradient of all the components of u , but, when multiplied by the gradient of the α component of u , from below we can see only the gradient of the α component of u . Componentwise coercivity allows us to arrive at Caccioppoli inequality on superlevel sets for the scalar function u^{α} , so that we can use De Giorgi's iteration for the scalar function u^{α} . Such a componentwise coercivity has been used by Bjorn (2001), Leonetti-Petricca (2011), Softova (2018), Palagacev-Softova (2020), Cupini-Leonetti-Mascolo (2022): [13], [93], [112], [111], [37].

Let us also mention Zhou (2000), [119], where a componentwise sign condition was assumed.

Examples of $A(x, z)$ enjoying componentwise coercivity are the Euler systems of some double phase functionals, see next section.

2. DOUBLE PHASE FUNCTIONALS

Example 1. Let us consider the variational integral

$$(12) \quad \mathcal{F}_1(u) = \int_{\Omega} (|Du(x)|^p + a(x)|Du(x)|^q) dx,$$

where

$$2 \leq p < q, \quad 0 \leq a(x) \leq M.$$

When $a(x) = 0$ then $|Du|^p + a(x)|Du|^q = |Du|^p$ and we are in the p -phase. On the other hand, if $a(x) > 0$ then, for large $|Du|$, we have that $|Du|^p + a(x)|Du|^q \approx |Du|^q$, so, we are

$$A_i^\alpha(x, z) = \frac{\partial f}{\partial z_i^\alpha}(x, z) = \frac{\partial g}{\partial t}(x, |z|) \frac{z_i^\alpha}{|z|},$$

so that

$$\sum_{i=1}^n A_i^\alpha(x, z) z_i^\alpha = \frac{\partial g}{\partial t}(x, |z|) \frac{|z^\alpha|^2}{|z|} \geq \nu |z|^{p-2} |z^\alpha|^2 \geq \nu |z^\alpha|^p,$$

so that componentwise coercivity holds true also for this more general structure.

REMARK. What happens to $f(x, z) = |z|^p + a(x)|z|^q$ when $1 < p < 2$? Componentwise coercivity fails in the vectorial case $m \geq 2$ when $1 < p < 2$. Let us consider this more general density

$$f(x, z) = (\mu + |z|^2)^{p/2} + a(x)|z|^q,$$

where $\mu \geq 0$, $a(x) = 0$ at some point x , $1 < p < 2$ and $m \geq 2$. The most important cases are $\mu = 0$, that gives $f(x, z) = |z|^p + a(x)|z|^q$, and $\mu = 1$, that gives $f(x, z) = (1 + |z|^2)^{p/2} + a(x)|z|^q$. Then

$$A_i^\alpha(x, z) = \frac{\partial f}{\partial z_i^\alpha}(x, z) = p(\mu + |z|^2)^{(p-2)/2} z_i^\alpha + a(x)q|z|^{q-2} z_i^\alpha.$$

We take x verifying $a(x) = 0$; then

$$A_i^\alpha(x, z) = p(\mu + |z|^2)^{(p-2)/2} z_i^\alpha$$

and

$$\sum_{i=1}^n A_i^\alpha(x, z) z_i^\alpha = p(\mu + |z|^2)^{(p-2)/2} |z^\alpha|^2.$$

By contradiction, let us assume that

$$(13) \quad \nu |z^\alpha|^{\tilde{p}} - c \leq \sum_{i=1}^n A_i^\alpha(x, z) z_i^\alpha$$

cut it!

for some positive constants ν, \tilde{p}, c . Since $m \geq 2$, then, for $\alpha \in \{1, \dots, m\}$, there exists $\beta \in \{1, \dots, m\} \setminus \{\alpha\}$. Let us take the matrix z as follows

$$z_1^\alpha = t > 0, \quad z_1^\beta = t^s, \quad z_i^\gamma = 0 \text{ otherwise,}$$

where $s > 1$ has to be chosen later. Then,

$$|z^\alpha| = t, \quad |z|^2 = t^2 + t^{2s} \quad (\mu + |z|^2)^{(p-2)/2} = (\mu + t^2 + t^{2s})^{(p-2)/2}.$$

Now, (13) becomes

$$\nu t^{\tilde{p}} - c \leq p(\mu + t^2 + t^{2s})^{(p-2)/2} t^2 = \frac{pt^2}{(\mu + t^2 + t^{2s})^{(2-p)/2}}.$$

We divide both sides by $t^{\tilde{p}}$ and we get

$$(14) \quad \nu - \frac{c}{t^{\tilde{p}}} \leq \frac{pt^{2-\tilde{p}}}{(\mu + t^2 + t^{2s})^{(2-p)/2}} = \frac{pt^{2-\tilde{p}}}{t^{s(2-p)} \left(\frac{\mu}{t^{2s}} + \frac{1}{t^{2s-2}} + 1 \right)^{(2-p)/2}}.$$

We would like that $s(2-p) > 2 - \tilde{p}$: since $p < 2$ we have $2-p > 0$ and this means that $s > \frac{2-\tilde{p}}{2-p}$. We choose $s > \max\{1; \frac{2-\tilde{p}}{2-p}\}$ and we let $t \rightarrow +\infty$: (14) becomes

$$\nu \leq 0.$$

This gives us the desired contradiction since ν was assumed to be positive: $\nu > 0$. The previous calculations were performed at points x with $a(x) = 0$. This means that also the standard vectorial p -Laplacian does not enjoy componentwise coercivity when $1 < p < 2$. More precisely, neither the system

$$\left\{ \begin{array}{l} -\sum_{i=1}^n D_i [p|Du|^{p-2} D_i u^1] = 0, \\ -\sum_{i=1}^n D_i [p|Du|^{p-2} D_i u^2] = 0, \\ \dots\dots\dots \\ -\sum_{i=1}^n D_i [p|Du|^{p-2} D_i u^m] = 0, \end{array} \right.$$

nor the one

$$\left\{ \begin{array}{l} -\sum_{i=1}^n D_i [p(1 + |Du|^2)^{(p-2)/2} D_i u^1] = 0, \\ -\sum_{i=1}^n D_i [p(1 + |Du|^2)^{(p-2)/2} D_i u^2] = 0, \\ \dots\dots\dots \\ -\sum_{i=1}^n D_i [p(1 + |Du|^2)^{(p-2)/2} D_i u^m] = 0, \end{array} \right.$$

enjoy the componentwise coercivity when $1 < p < 2$. Are there examples of subquadratic systems enjoying componentwise coercivity? Here is the following one in which we split the components:

$$(15) \quad \left\{ \begin{array}{l} - \sum_{i=1}^n D_i [p|Du^1|^{p-2}D_i u^1] = 0, \\ - \sum_{i=1}^n D_i [p|Du^2|^{p-2}D_i u^2] = 0, \\ \dots\dots\dots \\ - \sum_{i=1}^n D_i [p|Du^m|^{p-2}D_i u^m] = 0. \end{array} \right.$$

Indeed, in the previous system we have $A_i^\alpha(z) = p|z^\alpha|^{p-2}z_i^\alpha$ and componentwise coercivity can be easily checked as follows

$$\sum_{i=1}^n A_i^\alpha(z)z_i^\alpha = p|z^\alpha|^{p-2}|z^\alpha|^2 = p|z^\alpha|^p.$$

Note that the previous system is decoupled. The next one is no longer decoupled:

$$(16) \quad \left\{ \begin{array}{l} - \sum_{i=1}^n D_i [p|Du^1|^{p-2}D_i u^1] - D_1 \left[\frac{2D_1 u^1 D_1 u^2 D_1 u^2}{(1+(D_1 u^1 D_1 u^2)^2)^2} \right] = 0, \\ - \sum_{i=1}^n D_i [p|Du^2|^{p-2}D_i u^2] - D_1 \left[\frac{2D_1 u^1 D_1 u^2 D_1 u^1}{(1+(D_1 u^1 D_1 u^2)^2)^2} \right] = 0. \end{array} \right.$$

Here, $1 < p < 2$, $m = 2$ and $A_i^\alpha(z) = p|z^\alpha|^{p-2}z_i^\alpha + \frac{2z_1^1 z_1^2 z_1^{\hat{\alpha}}}{(1+(z_1^1 z_1^2)^2)^2}$, where $\hat{\alpha}$ is δ_{i1}

$$\hat{\alpha} = \begin{cases} 2 & \text{if } \alpha = 1, \\ 1 & \text{if } \alpha = 2. \end{cases}$$

Componentwise coercivity can be easily checked as follows

$$\sum_{i=1}^n A_i^\alpha(z)z_i^\alpha = p|z^\alpha|^{p-2}|z^\alpha|^2 + \frac{2z_1^1 z_1^2 z_1^{\hat{\alpha}} z_1^\alpha}{(1+(z_1^1 z_1^2)^2)^2} = p|z^\alpha|^p + \frac{2(z_1^1 z_1^2)^2}{(1+(z_1^1 z_1^2)^2)^2} \geq p|z^\alpha|^p.$$

On the other hand, system (16) does not enjoy Uhlenbeck structure. Indeed, we argue by contradiction: let assume Uhlenbeck structure, so that

$$(17) \quad A_i^\alpha(z) = g'(|z|) \frac{z_i^\alpha}{|z|}.$$

We first take the matrix \tilde{z} as follows

$$\tilde{z}_1^1 = 1, \quad \tilde{z}_j^\beta = 0 \text{ otherwise,}$$

so that $|\tilde{z}| = |\tilde{z}^1| = 1$. Then, we take the matrix $\tilde{\tilde{z}}$ as follows

$$\tilde{\tilde{z}}_1^1 = \frac{1}{\sqrt{2}}, \quad \tilde{\tilde{z}}_2^2 = \frac{1}{\sqrt{2}}, \quad \tilde{\tilde{z}}_j^\beta = 0 \text{ otherwise,}$$

so that $|\tilde{\tilde{z}}| = 1$ and $|\tilde{\tilde{z}}^1| = \frac{1}{\sqrt{2}}$. We first exploit (17) with $z = \tilde{z}$:

$$A_1^1(\tilde{z}) = g'(1).$$

Then, we exploit (17) with $z = \tilde{\tilde{z}}$:

$$A_1^1(\tilde{\tilde{z}}) = g'(1) \frac{1}{\sqrt{2}}.$$

These two conditions merge into

$$(18) \quad A_1^1(\tilde{\tilde{z}}) = g'(1) \frac{1}{\sqrt{2}} = A_1^1(\tilde{z}) \frac{1}{\sqrt{2}}.$$

Now, we compute $A_1^1(\tilde{\tilde{z}})$ and $A_1^1(\tilde{z})$. We have $A_1^1(\tilde{\tilde{z}}) = p \left(\frac{1}{\sqrt{2}}\right)^{p-1}$ and $A_1^1(\tilde{z}) = p$. Then, (18) gives

$$p \left(\frac{1}{\sqrt{2}}\right)^{p-1} = p \frac{1}{\sqrt{2}},$$

that is $2^{\frac{1-p}{2}} = 2^{\frac{-1}{2}}$; this means $\frac{1-p}{2} = \frac{-1}{2}$, that is, $2 = p$: this is false, since we assumed $1 < p < 2$. This shows that we cannot have Uhlenbeck structure in system (16).

Functional \mathcal{F}_2 is inspired by an example contained in [80], Hasto-Ok (2022). Let us consider the density of \mathcal{F}_2 :

$$f(x, z) = |z|^p + a(x) \sum_{\beta=1}^m \sum_{j=1}^n (z_j^\beta)^4,$$

so that

$$A_i^\alpha(x, z) = \frac{\partial f}{\partial z_i^\alpha}(x, z) = p|z|^{p-2}z_i^\alpha + a(x)4(z_i^\alpha)^3.$$

Componentwise coercivity holds true: see [91]. On the contrary, Uhlenbeck structure does not hold true, see [91]. The Euler system of functional \mathcal{F}_2 appears to be as follows

$$\left\{ \begin{array}{l} -\sum_{i=1}^n D_i [p|Du|^{p-2}D_i u^1 + a(x)4(D_i u^1)^3] = 0, \\ -\sum_{i=1}^n D_i [p|Du|^{p-2}D_i u^2 + a(x)4(D_i u^2)^3] = 0, \\ \dots\dots\dots \\ -\sum_{i=1}^n D_i [p|Du|^{p-2}D_i u^m + a(x)4(D_i u^m)^3] = 0. \end{array} \right.$$

Example 3. Let us consider the variational integral

$$(21) \quad \mathcal{F}_3(u) = \int_{\Omega} \{ |Du(x)|^p + a(x) (\max\{D_n u^1(x); 0\})^q \} dx,$$

where

$$2 \leq p < q, \quad 0 \leq a(x) \leq M.$$

Functional \mathcal{F}_3 is inspired by an example in [117], Tang (1993). See also Esposito-Leonetti-Petricca (2019): [61].

Let us consider the density of \mathcal{F}_3 :

$$f(x, z) = |z|^p + a(x) (\max\{z_n^1; 0\})^q,$$

so that

$$A_i^\alpha(x, z) = \frac{\partial f}{\partial z_i^\alpha}(x, z) = p|z|^{p-2}z_i^\alpha + a(x) \delta^{\alpha 1} \delta_{in} q (\max\{z_n^1; 0\})^{q-1},$$

where δ_{in} is Kronecker symbol: $\delta_{in} = 1$ if $i = n$ and $\delta_{in} = 0$ otherwise; in a similar manner $\delta^{\alpha 1}$ is defined. Componentwise coercivity holds true: see [91]. Does this example enjoy Uhlenbeck structure? No, see [91]. Euler system of \mathcal{F}_3 appears to be

$$\left\{ \begin{array}{l} -\sum_{i=1}^n D_i [p|Du|^{p-2}D_i u^1] - D_n [a(x)q (\max\{D_n u^1; 0\})^{q-1}] = 0, \\ -\sum_{i=1}^n D_i [p|Du|^{p-2}D_i u^2] = 0, \\ \dots\dots\dots \\ -\sum_{i=1}^n D_i [p|Du|^{p-2}D_i u^m] = 0. \end{array} \right.$$

Example 4. Let us consider the variational integral

$$(22) \quad \mathcal{F}_4(u) = \int_{\Omega} \left\{ |Du(x)|^p + a(x) (D_1 u^1(x) - D_2 u^1(x))^4 \right\} dx,$$

where

$$2 \leq p < 4 = q, \quad 0 \leq a(x) \leq M.$$

This functional has been inspired by an example in [31], Cianchi (2000); see also Esposito-Leonetti-Petricca (2019): [61].

Let us consider the density of \mathcal{F}_4 :

$$f(x, z) = |z|^p + a(x) (z_1^1 - z_2^1)^4;$$

then

$$A_i^\alpha(x, z) = \frac{\partial f}{\partial z_i^\alpha}(x, z) = p|z|^{p-2}z_i^\alpha + a(x) \delta^{\alpha 1} 4 (z_1^1 - z_2^1)^{4-1} [\delta_{i1} - \delta_{i2}].$$

$$\nu(\mu + |z|^2 + |\tilde{z}|^2)^{\frac{p-2}{2}}|z - \tilde{z}|^2 \leq \left\langle \frac{\partial f}{\partial z}(x, z) - \frac{\partial f}{\partial z}(x, \tilde{z}); z - \tilde{z} \right\rangle$$

and

$$\left| \frac{\partial f}{\partial z}(x, z) - \frac{\partial f}{\partial z}(y, z) \right| \leq c|x - y|^\sigma(1 + |z|^{q-1}).$$

Moreover, we assume that

$$1 < p < q < p \frac{n + \sigma}{n},$$

and

$$\mathcal{F}(u_k, B_R) \rightarrow \mathcal{F}(u, B_R)$$

for some $u_k \in W^{1,p}(B_R; \mathbb{R}^m) \cap W_{loc}^{1,q}(B_R; \mathbb{R}^m)$ with $u_k \rightarrow u$ weakly in $W^{1,p}(B_R; \mathbb{R}^m)$.

Under these assumptions, the minimizer u enjoys the following higher integrability of the gradient

$$u \in W_{loc}^{1,q}(\Omega; \mathbb{R}^m).$$

This is contained in Esposito-Leonetti-Mingione (2004): [64].

Failure of the energy approximation

$$\mathcal{F}(u_k, B_R) \rightarrow \mathcal{F}(u, B_R)$$

for some $u_k \in W^{1,p}(B_R; \mathbb{R}^m) \cap W_{loc}^{1,q}(B_R; \mathbb{R}^m)$ with $u_k \rightarrow u$ weakly in $W^{1,p}(B_R; \mathbb{R}^m)$, gives rise to the Lavrentiev phenomenon:

$$\inf_{v \in u + W_0^{1,p}(B_R; \mathbb{R}^m)} \mathcal{F}(v, B_R) < \inf_{v \in [u + W_0^{1,p}(B_R; \mathbb{R}^m)] \cap W_{loc}^{1,q}(B_R; \mathbb{R}^m)} \mathcal{F}(v, B_R),$$

see the survey [10] and the recent contributions [118], [44], [66], [64], [45], [79], [19], [6], [18], [4].

Do the double phase functionals $\mathcal{F}_1(u, \Omega), \dots, \mathcal{F}_4(u, \Omega)$ enjoy the energy approximation?
YES!

$$\begin{aligned}\mathcal{F}_1(u, \Omega) &= \int_{\Omega} (|Du(x)|^p + a(x)|Du(x)|^q) dx, \\ \mathcal{F}_2(u, \Omega) &= \int_{\Omega} \left\{ |Du(x)|^p + a(x) \sum_{\beta=1}^m \sum_{j=1}^n (D_j u^\beta(x))^4 \right\} dx, \\ \mathcal{F}_3(u, \Omega) &= \int_{\Omega} \{ |Du(x)|^p + a(x) (\max\{D_n u^1(x); 0\})^q \} dx, \\ \mathcal{F}_4(u) &= \int_{\Omega} \left\{ |Du(x)|^p + a(x) (D_1 u^1(x) - D_2 u^1(x))^4 \right\} dx,\end{aligned}$$

$$a \in C^{0,\sigma}(\Omega), \quad q \leq p \frac{n + \sigma}{n},$$

↓

$$\mathcal{F}_i(u_k, B_R) \rightarrow \mathcal{F}_i(u, B_R)$$

for some $u_k \in W^{1,p}(B_R; \mathbb{R}^m) \cap W_{loc}^{1,q}(B_R; \mathbb{R}^m)$ with $u_k \rightarrow u$ weakly in $W^{1,p}(B_R; \mathbb{R}^m)$. For energy approximation, see [118], [64], [61], [47], [51], [86], [24], [53], [18], [50], [54].

Remark. Existence of $W^{1,q}$ solutions to general elliptic systems with p, q -growth is obtained in [35]. Existence of solutions to one single equation with p, q -growth and with explicit dependence on u has been studied in the recent [41].

An interesting example

Example 5. Let us consider the variational integral

$$(23) \quad \mathcal{F}_5(u) = \int_{\Omega} \{ |Du(x) - \hat{z}|^p + a(x)|Du(x)|^q \} dx,$$

where

$$\hat{z} \in \mathbb{R}^{m \times n}, \quad \hat{z} \neq 0, \quad 2 \leq p < q, \quad 0 \leq a(x) \leq M.$$

This functional has been inspired by an example in [76], Guarnotta-Mosconi (2023). Let us consider the density of \mathcal{F}_5 :

$$f(x, z) = |z - \hat{z}|^p + a(x)|z|^q,$$

so that

$$A_i^\alpha(x, z) = \frac{\partial f}{\partial z_i^\alpha}(x, z) = p|z - \hat{z}|^{p-2}(z_i^\alpha - \hat{z}_i^\alpha) + a(x)q|z|^{q-2}z_i^\alpha.$$

Does this example satisfy Uhlenbeck structure? No, see [91]. What about componentwise coercivity? The answer depends on p :

$$2 = p \quad \implies \quad \text{Yes}$$

$$2 < p \quad \implies \quad \text{No}$$

Let us check componentwise coercivity when $2 = p$. We write the density for \mathcal{F}_5 when $2 = p$:

$$f(x, z) = |z - \hat{z}|^2 + a(x)|z|^q,$$

so that

$$A_i^\alpha(x, z) = \frac{\partial f}{\partial z_i^\alpha}(x, z) = 2(z_i^\alpha - \hat{z}_i^\alpha) + a(x)q|z|^{q-2}z_i^\alpha.$$

Then

$$\sum_{i=1}^n A_i^\alpha(x, z)z_i^\alpha = 2\langle z^\alpha - \hat{z}^\alpha; z^\alpha \rangle + a(x)q|z|^{q-2}|z^\alpha|^2 \geq$$

$$2\langle z^\alpha - \hat{z}^\alpha; z^\alpha \rangle = 2|z^\alpha|^2 - 2\langle \hat{z}^\alpha; z^\alpha \rangle \geq 2|z^\alpha|^2 - 2|\hat{z}^\alpha||z^\alpha| \geq$$

$$2|z^\alpha|^2 - |\hat{z}^\alpha|^2 - |z^\alpha|^2 = |z^\alpha|^2 - |\hat{z}^\alpha|^2,$$

REFERENCES

- [1] E. Acerbi, N. Fusco. *Partial regularity under anisotropic (p,q) growth conditions*, J. Differential Equations **107** (1994), no. 1, 46-67.
- [2] A. Alberico, I. Chlebicka, A. Cianchi, A. Zatorska-Goldstein. *Fully anisotropic elliptic problems with minimally integrable data* Calc. Var. Partial Differential Equations **58** (2019), no. 6, Paper No. 186, 50 pp.
- [3] A. Alberico, G. di Blasio, F. Feo. *Estimates for fully anisotropic elliptic equations with a zero order term* Nonlinear Anal. **181** (2019), 249-264.
- [4] S. Almi, S. Kromer, A. Molchanova. *A new example for the Lavrentiev phenomenon in nonlinear elasticity* Z. Angew. Math. Phys. **75** (2024), no.1, Paper No. 2, 21 pp.
- [5] E. Amoroso, A. Sciammetta, P. Winkert. *Anisotropic (p,q) -Laplacian problems with superlinear nonlinearities* Commun. Anal. Mech. **16** (2024), no.1, 1-23.
- [6] A. K. Balci, L. Diening, M. Surnachev. *New examples on Lavrentiev gap using fractals* Calc. Var. Partial Differential Equations **59** (2020), no. 5, Paper No. 180, 34 pp.
- [7] P. Baroni, M. Colombo, G. Mingione. *Harnack inequalities for double phase functionals* Nonlinear Anal. **121** (2015), 206-222.
- [8] L. Beck, G. Mingione. *Lipschitz bounds and nonuniform ellipticity* Comm. Pure Appl. Math. **73** (2020), no. 5, 944-1034.
- [9] P. Bella, M. Schäffner. *On the regularity of minimizers for scalar integral functionals with (p,q) -growth* Anal. PDE **13** (2020), no. 7, 2241-2257.
- [10] M. Belloni, G. Buttazzo. *A survey of old and recent results about the gap phenomenon in the calculus of variations* in: Recent Developements in Well-Posed Variational Problems, Math. Appl. **331** Kluwer Academic, Dordrecht (1995), 1-27.
- [11] T. Bhattacharya, F. Leonetti. *$W^{2,2}$ regularity for weak solutions of elliptic systems with nonstandard growth* J. Math. Anal. Appl. **176** (1993), no. 1, 224-234.
- [12] M. Bildhauer, M. Fuchs. *$C^{1,\alpha}$ -solutions to non-autonomous anisotropic variational problems* Calc. Var. Partial Differential Equations **24** (2005), no. 3, 309-340.
- [13] J. Bjorn. *Boundedness and differentiability for nonlinear elliptic systems* Trans. Amer. Math. Soc. **353** (2001), no. 11, 4545-4565.
- [14] L. Boccardo, P. Marcellini, C. Sbordone. *L^∞ -regularity for variational problems with sharp non-standard growth conditions* Boll. Un. Mat. Ital. A (7) **4** (1990), no. 2, 219-225.
- [15] V. Bogelein, F. Duzaar, J. Habermann, C. Scheven. *Stationary electro-rheological fluids: low order regularity for systems with discontinuous coefficients* Adv. Calc. Var. **5** (2012), no.1, 1-57.

- [16] V. Bogelein, F. Duzaar, P. Marcellini, C. Scheven. *Boundary regularity for elliptic systems with p, q -growth* J. Math. Pures Appl. (9) **159** (2022), 250-293.
- [17] G. Bonanno, G. D'Aguí, A. Sciammetta. *Multiple solutions for a class of anisotropic p -Laplacian problems* Bound. Value Probl. (2023), Paper No. 89, 12 pp.
- [18] M. Borowski, I. Chlebicka, F. De Filippis, B. Miasojedow. *Absence and presence of Lavrentiev's phenomenon for double phase functionals upon every choice of exponents* Calc. Var. Partial Differential Equations **63** (2024), no. 2, Paper No. 35, 23 pp.
- [19] P. Bousquet, C. Mariconda, G. Treu. *A survey on the non occurrence of the Lavrentiev gap for convex, autonomous multiple integral scalar variational problems* Set-Valued Var. Anal. **23** (2015), no.1, 55-68.
- [20] P. Bousquet, L. Brasco, C. Leone. *Singular orthotropic functionals with nonstandard growth conditions* Rev. Mat. Iberoam. **40** (2024), no.2, 753-802.
- [21] L. Brasco, C. Leone, G. Pisante, A. Verde. *Sobolev and Lipschitz regularity for local minimizers of widely degenerate anisotropic functionals* Nonlinear Anal. **153** (2017), 169-199.
- [22] F. Brock, J. I. Diaz, A. Ferone, D. Gomez-Castro, A. Mercaldo. *Steiner symmetrization for anisotropic quasilinear equations via partial discretization* Ann. Inst. H. Poincaré Anal. Non Linéaire **38** (2021), no.2, 347-368.
- [23] M. Bulíček, G. Cupini, B. Stroffolini, A. Verde. *Existence and regularity results for weak solutions to (p, q) -elliptic systems in divergence form* Adv. Calc. Var. **11** (2018), no.3, 273-288.
- [24] M. Bulíček, P. Gwiazda, J. Skrzeczkowski. *On a Range of Exponents for Absence of Lavrentiev Phenomenon for Double Phase Functionals* Arch. Rational Mech. Anal. **246** (2022) 209-240.
- [25] A. Canale, A. D'Ottavio, F. Leonetti, M. Longobardi. *Differentiability for bounded minimizers of some anisotropic integrals* J. Math. Anal. Appl. **253** (2001), no. 2, 640-650.
- [26] M. Carozza, J. Kristensen, A. Passarelli di Napoli. *Regularity of minimizers of autonomous convex variational integrals* Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **13** (2014), no.4, 1065-1089.
- [27] M. Carozza, H. Gao, R. Giova, F. Leonetti. *A boundedness result for minimizers of some polyconvex integrals* J. Optim. Theory Appl. **178** (2018), no. 3, 699-725.
- [28] P. Cavaliere, A. D'Ottavio, F. Leonetti, M. Longobardi. *Differentiability for minimizers of anisotropic integrals* Comment. Math. Univ. Carolin. **39** (1998), no. 4, 685-696.
- [29] H. J. Choe. *Regularity for minimizers of certain degenerate functionals with nonstandard growth conditions* Comm. Partial Differential Equations **16** (1991), no. 2-3, 363-372.
- [30] H. J. Choe. *Interior behaviour of minimizers for certain functionals with nonstandard growth* Nonlinear Anal. **19** (1992), no. 10, 933-945.
- [31] A. Cianchi. *Local boundedness of minimizers of anisotropic functionals* Ann. Inst. H. Poincaré Anal. Non Linéaire **17** (2000), no. 2, 147-168.

- [32] A. Cianchi. *Symmetrization in anisotropic elliptic problems* Comm. Partial Differential Equations **32** (2007), no.4-6, 693-717.
- [33] G. Cupini, M. Focardi, F. Leonetti, E. Mascolo. *On the Hölder continuity for a class of vectorial problems* Adv. Nonlinear Anal. **9** (2020), no. 1, 1008-1025.
- [34] G. Cupini, F. Giannetti, R. Giova, A. Passarelli di Napoli. *Regularity results for vectorial minimizers of a class of degenerate convex integrals* J. Differential Equations **265** (2018), no. 9, 4375-4416.
- [35] G. Cupini, F. Leonetti, E. Mascolo. *Existence of weak solutions for elliptic systems with p, q -growth* Ann. Acad. Sci. Fenn. Math. **40** (2015), no. 2, 645-658.
- [36] G. Cupini, F. Leonetti, E. Mascolo. *Local boundedness for minimizers of some polyconvex integrals* Arch. Ration. Mech. Anal. **224** (2017) 269-289.
- [37] G. Cupini, F. Leonetti, E. Mascolo. *Local boundedness for solutions of a class of nonlinear elliptic systems* Calc. Var. Partial Differential Equations **61** (2022), no. 3, Paper No. 94, 17 pp.
- [38] G. Cupini, P. Marcellini, E. Mascolo. *Local boundedness of solutions to quasilinear elliptic systems* Manuscripta Math. **137** (2012) 287-315.
- [39] G. Cupini, P. Marcellini, E. Mascolo. *Local boundedness of solutions to some anisotropic elliptic systems* Recent trends in nonlinear partial differential equations. II. Stationary problems 169-186, Contemp. Math., 595, Amer. Math. Soc., Providence, RI, 2013.
- [40] G. Cupini, P. Marcellini, E. Mascolo. *Regularity for nonuniformly elliptic equations with p, q -growth and explicit x, u -dependence* Arch. Ration. Mech. Anal. **248** 60 (2024). <https://doi.org/10.1007/s00205-024-01982-0>
- [41] G. Cupini, P. Marcellini, E. Mascolo. *The Leray-Lions existence theorem under general growth conditions* <https://arxiv.org/abs/2309.15811>
- [42] G. Cupini, E. Mascolo. *Local regularity for elliptic systems with p, q -growth* Bruno Pini Math. Anal. Semin. 2015 Università di Bologna, Alma Mater Studiorum, Bologna, 2015, 15-38.
- [43] A. Dall'Aglio, E. Mascolo, G. Papi. *Local boundedness for minima of functionals with nonstandard growth conditions* Rend. Mat. Appl. (7) **18** (1998), no.2, 305-326.
- [44] R. De Arcangelis, C. Trombetti. *On the Lavrentieff phenomenon for some classes of Dirichlet minimum points* J. Convex Anal. **7** (2000), no.2, 271-297.
- [45] R. De Arcangelis, S. Monsurró, E. Zappale. *On the relaxation and the Lavrentieff phenomenon for variational integrals with pointwise measurable gradient constraints* Calc. Var. Partial Differential Equations **21** (2004), no. 4, 357-400.
- [46] C. De Filippis, F. Leonetti. *Uniform ellipticity and (p, q) growth* J. Math. Anal. Appl. **501** (2021), no. 1, Paper No. 124451, 11 pp.
- [47] C. De Filippis, G. Mingione. *On the regularity of minima of non-autonomous functionals* J. Geom. Anal. **30** (2020), no. 2, 1584-1626.

- [48] C. De Filippis, G. Mingione. *Lipschitz bounds and nonautonomous integrals* Arch. Ration. Mech. Anal. **242** (2021), no. 2, 973-1057.
- [49] C. De Filippis, G. Mingione. *Nonuniformly elliptic Schauder theory* Invent. math. **234** (2023) 1109-1196.
- [50] C. De Filippis, G. Mingione. *Regularity for Double Phase Problems at Nearly Linear Growth* Arch. Ration. Mech. Anal. **247** 85 (2023).
- [51] F. De Filippis, F. Leonetti. *No Lavrentiev gap for some double phase integrals* Adv. Calc. Var. **17** (2024), no. 1, 165-194.
- [52] F. De Filippis, F. Leonetti, P. Marcellini, E. Mascolo. *The Sobolev class where a weak solution is a local minimizer* Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl. **34** (2023), no. 2, 451-463.
- [53] F. De Filippis, F. Leonetti, G. Treu. *Non occurrence of Lavrentiev Gap for a class of functionals with non standard growth* Adv. Nonlinear Anal. **13** (2024). <https://doi.org/10.1515/anona-2024-0002>
- [54] F. De Filippis, M. Piccinini. *Regularity for multi-phase problems at nearly linear growth* J. Differential Equations **410** (2024), 832-868.
- [55] E. De Giorgi. *Sulla differenziabilità e l'analiticità delle estremali degli integrali multipli regolari* Mem. Accad. Sci. Torino. Cl. Sci. Fis. Mat. Nat. **3** (1957) 25-43.
- [56] E. De Giorgi. *Un esempio di estremali discontinue per un problema variazionale di tipo ellittico* Boll. Un. Mat. Ital. (4) **1** (1968) 135-137.
- [57] E. Di Benedetto, U. Gianazza, V. Vespri. *Remarks on local boundedness and local Hölder continuity of local weak solutions to anisotropic p -Laplacian type equations* J. Elliptic Parabol. Equ. **2** (2016), no.1-2, 157-169.
- [58] T. Di Marco, P. Marcellini. *A-priori gradient bound for elliptic systems under either slow or fast growth conditions* Calc. Var. Partial Differential Equations **59** (2020), no 4, Paper No. 120, 26 pp.
- [59] M. Eleuteri, P. Marcellini, E. Mascolo. *Regularity for scalar integrals without structure conditions* Adv. Calc. Var. **13** (2020), no.3, 279-300.
- [60] M. Eleuteri, P. Marcellini, E. Mascolo, S. Perrotta. *Local Lipschitz continuity for energy integrals with slow growth* Ann. Mat. Pura Appl. (4) **201** (2022), no.3, 1005-1032.
- [61] A. Esposito, F. Leonetti, P. V. Petricca. *Absence of Lavrentiev gap for non-autonomous functionals with (p,q) -growth* Adv. Nonlinear Anal. **8** (2019), 73-78.
- [62] L. Esposito, F. Leonetti, G. Mingione. *Regularity for minimizers of functionals with $p - q$ growth* NoDEA Nonlinear Differential Equations Appl. **6** (1999), no. 2, 133-148.
- [63] L. Esposito, F. Leonetti, G. Mingione. *Higher integrability for minimizers of integral functionals with (p,q) growth* J. Differential Equations **157** (1999), no. 2, 414-438.
- [64] L. Esposito, F. Leonetti, G. Mingione. *Sharp regularity for functionals with (p, q) -growth* J. Differential Equations **204** (2004), no. 1, 5-55.

- [65] I. Fonseca, J. Maly, G. Mingione. *Scalar minimizers with fractal singular sets* Arch. Ration. Mech. Anal. **172** (2004), 295-307.
- [66] M. Foss, W. J. Hrusa, V. J. Mizel. *The Lavrentiev gap phenomenon in nonlinear elasticity* Arch. Ration. Mech. Anal. **167** (2003), no.4, 337-365.
- [67] J. Frehse. *Una generalizzazione di un controesempio di De Giorgi nella teoria delle equazioni ellittiche* Boll. Un. Mat. Ital. (4) **3** (1970) 998-1002.
- [68] J. Frehse. *An irregular complex valued solution to a scalar uniformly elliptic equation* Calc. Var. Partial Differential Equations **33** (2008) 263-266.
- [69] N. Fusco, C. Sbordone. *Higher integrability of the gradient of minimizers of functionals with non-standard growth conditions* Comm. Pure Appl. Math. **43** (1990), no. 5, 673-683.
- [70] N. Fusco, C. Sbordone. *Some remarks on the regularity of minima of anisotropic integrals* Comm. P. D. E., **18** (1993) 153-167.
- [71] D. Giachetti, M. M. Porzio. *Local regularity results for minima of functionals of the calculus of variation* Nonlinear Anal. **39** (2000), no. 4, 463-482.
- [72] M. Giaquinta. *A counter-example to the boundary regularity of solutions to elliptic quasilinear systems* Manuscripta Math. **24** (1978), no. 2, 217-220.
- [73] M. Giaquinta. *Growth conditions and regularity, a counterexample* Manuscripta Math. **59** (1987) 245-248.
- [74] M. Giaquinta, J. Soucek. *Caccioppoli's inequality and Legendre-Hadamard condition* Math. Ann. **270** (1985), no.1, 105-107.
- [75] E. Giusti, M. Miranda. *Un esempio di soluzioni discontinue per un problema di minimo relativo ad un integrale regolare del calcolo delle variazioni* Boll. Un. Mat. Ital. **2** (1968) 1-8.
- [76] U. Guarnotta, S. Mosconi. *A general notion of uniform ellipticity and the regularity of the stress field for elliptic equations in divergence form* Anal. PDE **16** (2023), no. 8, 1955-1988.
- [77] W. Hao, S. Leonardi, J. Necas. *An example of irregular solution to a nonlinear Euler-Lagrange elliptic system with real analytic coefficients* Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **23** (1996), no. 1, 57-67.
- [78] W. Hao, S. Leonardi, M. Steinhauer. *Examples of discontinuous, divergence-free solutions to elliptic variational problems* Comment. Math. Univ. Carolin. **36** (1995), no. 3, 511-517.
- [79] P. A. Hästö. *Counter-examples of regularity in variable exponent Sobolev spaces* The p-harmonic equation and recent advances in analysis, 133-143. Contemp. Math., 370 American Mathematical Society, Providence, RI, 2005
- [80] P. A. Hästö, J. Ok. *Regularity Theory for Non-autonomous Partial Differential Equations Without Uhlenbeck Structure* Arch. Ration. Mech. Anal. **245** (2022), 1401-1436.

- [81] J. Hirsch, M. Scäffner. *Growth conditions and regularity, an optimal local boundedness result* Commun. Contemp. Math. **23** (2021), no. 3, Paper No. 2050029, 17 pp.
- [82] M. C. Hong. *Some remarks on the minimizers of variational integrals with nonstandard growth conditions* Boll. Un. Mat. Ital. A (7) **6** (1992), no.1, 91-101.
- [83] C. Irving, L. Koch. *Boundary regularity results for minimisers of convex functionals with (p,q) -growth* Adv. Nonlinear Anal. **12** (2023), no. 1, Paper No. 20230110, 57 pp.
- [84] O. John, J. Maly, J. Stara. *Nowhere continuous solutions to elliptic systems* Comment. Math. Univ. Carolin. **30** (1989), no. 1, 33-43.
- [85] O. John, J. Necas, J. Stara. *Counterexample to the regularity of weak solution of elliptic systems* Comment. Math. Univ. Carolin. **21** (1980), no. 1, 145-154.
- [86] L. Kock. *Global higher integrability for minimisers of convex functionals with (p,q) -growth* Calc. Var. Partial Differential Equations **60** (2021), no. 2, Paper No. 63.
- [87] J. Kristensen, G. Mingione. *Sketches of Regularity Theory from The 20th Century and the Work of Jindrich Necas* in “Selected works of Jindrich Necas”. PDEs, continuum mechanics and regularity. Edited by Sarka Necasova, Milan Pokorny and Vladimir Sverak. Advances in Mathematical Fluid Mechanics. *Birkhauser/Springer, Basel*, 2015. xiv+785 pp.
- [88] R. Landes. *Some remarks on bounded and unbounded weak solutions of elliptic systems* Manuscripta Math. **64** (1989) 227-234.
- [89] M. Lavrentiev. *Sur quelques problemes du calcul des variations* Ann. Math. Pura Appl. **4** (1926) 107-124.
- [90] F. Leonetti. *Regularity results for minimizers of integral functionals with nonstandard growth* Atti Sem. Mat. Fis. Univ. Modena **43** (1995), no. 2, 425-429.
- [91] F. Leonetti. *Elliptic systems and double phase functionals* Atti Sem. Mat. Fis. Univ. Modena (to appear).
- [92] F. Leonetti, P. V. Petricca. *Regularity for vector valued minimizers of some anisotropic integral functionals* JIPAM. J. Inequal. Pure Appl. Math. **7** (2006), no. 3, Article 88, 7 pp.
- [93] F. Leonetti, P.V. Petricca. *Regularity for solutions to some nonlinear elliptic systems* Complex Var. Elliptic Equ. **56** (2011), no. 12, 1099-1113.
- [94] G. M. Lieberman. *The natural generalization of the natural conditions of Ladyzhenskaya and Ural'tseva for elliptic equations* Comm. Partial Differential Equations **16** (1991), no. 2-3, 311-361.
- [95] B. Maniá. *Sopra un esempio di Lavrentieff* Boll. Un. Mat. Ital. **13** (1934), 147-153.
- [96] P. Marcellini. *Un esempio de solution discontinue d'un problème variationnel dans le cas scalaire* Preprint 11, Istituto Matematico “U.Dini”, Università di Firenze, 1987.
- [97] P. Marcellini. *Regularity of minimizers of integrals of the calculus of variations with nonstandard growth conditions* Arch. Rational Mech. Anal. **105** (1989), no. 3, 267-284.

- [98] P. Marcellini. *Regularity and existence of solutions of elliptic equations with p, q -growth conditions* J. Differential Equations **90** (1991), no. 1, 1-30.
- [99] P. Marcellini. *Regularity under general and p, q -growth conditions* Discrete Cont. Dynamical Systems Series S **13** (2020), 2009-2031.
- [100] P. Marcellini. *Growth conditions and regularity for weak solutions to nonlinear elliptic pdes* J. Math. Anal. Appl. **501** (2021), no. 1, 124408, 32 pp.
- [101] V. G. Maz'ja. *Examples of nonregular solutions of quasilinear elliptic equations with analytic coefficients* (Russian) Funkcional. Anal. i Priložen bf 2 (1968) no. 3, 53-57.
- [102] G. Mingione. *Regularity of minima: an invitation to the dark side of the calculus of variations* Appl. Math. **51** (2006) 355-426.
- [103] G. Mingione. *Singularities of minima: a walk on the wild side of the calculus of variations* J. Global Optim. **40** (2008) 209-223.
- [104] G. Mingione, V. Radulescu. *Recent developments in problems with nonstandard growth and nonuniform ellipticity* J. Math. Anal. Appl. **501** (2021), no. 1, 125197, 41 pp.
- [105] C. Mooney, O. Savin. *Some singular minimizers in low dimensions in the calculus of variations* Arch. Ration. Mech. Anal. **221** (2016), 1-22.
- [106] G. MoscarIELLO, L. Nania. *Hölder continuity of minimizers of functionals with non standard growth conditions* Ricerche Mat. **40** (1991), 259-273.
- [107] J. Moser. *A new proof of De Giorgi's theorem concerning the regularity problem for elliptic differential equations* Commun. Pure Appl. Math. **13** (1960), 457-468.
- [108] J. Nash. *Continuity of solutions of parabolic and elliptic equations* Am. J. Math. **80** (1958), 931-954.
- [109] A. Nastasi, C. Pacchiano Camacho. *Higher integrability and stability of (p, q) -quasiminimizers* J. Differential Equations **342** (2023), 121-149.
- [110] J. Nečas. *Example of an irregular solution to a nonlinear elliptic system with analytic coefficients and conditions for regularity* Theory Nonlin. Oper. Abhand. der Wiss. der DDR (1977).
- [111] D. Palagachev, L. Softova. *Boundedness of solutions to a class of coercive systems with Morrey data* Nonlinear Anal. **191** (2020), 111630, 16 pp.
- [112] L. Softova. *Boundedness of the solutions to non-linear systems with Morrey data* Complex Var. Elliptic Equ. **63** (2018), no. 11, 1581-1594.
- [113] J. Soucek. *Singular solutions to linear elliptic systems* Comment. Math. Univ. Carolin. **25** (1984), no. 2, 273-281.
- [114] B. Stroffolini. *Global boundedness of solutions of anisotropic variational problems* Boll. Un. Mat. Ital. A (7) **5** (1991), no. 3, 345-352.
- [115] V. Sverák, X. Yan. *A singular minimizer of a smooth strongly convex functional in three dimensions* Calc. Var. Partial Differential Equations **10** no. 3 (2000) 213-221.

- [116] G. Talenti. *Boundedness of minimizers* Hokkaido Math. J. **19** (1990), no.2, 259-279.
- [117] Q. Tang. *Regularity of minimizers of nonisotropic integrals in the calculus of variations* Ann. Mat. Pura Appl. (4) **164** (1993), 77-87.
- [118] V. V. Zhikov. *On Laurentiev's phenomenon* Russ. J. Math. Phys. **3** (1995), 249-269.
- [119] S. Zhou. *A note on nonlinear elliptic systems involving measures* Electron J. Differ. Equ. 2000 (08) (2000) 1-6.

UNIVERSITÀ DEGLI STUDI DELL'AQUILA, DISIM, VIA VETOIO SNC, COPPITO, 67100 – L'AQUILA,
ITALY

Email address: francesco.leonetti@univaq.it