

Article

Dissipation Functions and Brownian Oscillators

Matteo Colangeli ^{1,*} , Lamberto Rondoni ^{2,3}  and Pasquale Vozza ⁴ 

¹ Dipartimento di Ingegneria e Scienze dell'Informazione e Matematica, Università degli Studi dell'Aquila, Via Vetoio, 67100 L'Aquila, Italy

² Dipartimento di Scienze Matematiche, Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino, Italy; lamberto.rondoni@polito.it

³ Istituto Nazionale di Fisica Nucleare, Sezione di Torino, Via Pietro Giuria 1, 10125 Torino, Italy

⁴ Dipartimento di Fisica, Università degli Studi di Torino, Via Pietro Giuria 1, 10125 Torino, Italy; pasquale.vozza@edu.unito.it

* Correspondence: matteo.colangeli1@univaq.it

Abstract

We develop a general framework for response theory in Markovian diffusion processes governed by Fokker–Planck equations. Our formalism, based on the notion of the dissipation function, is capable of handling dynamics that are driven by an external perturbation arbitrarily far from a given reference process. Using the analytically solvable Brownian oscillator model, we derive exact response formulae for both overdamped and underdamped dynamics of harmonically bound Brownian particles. We also demonstrate that for certain observables and under suitable time scaling, the operations of model reduction and response formula extraction commute, which highlights a relevant symmetry of the adopted mathematical formalism. Notably, the time reversal invariance symmetry, which is manifested as detailed balance in stochastic processes and is often required in statistical mechanics, is not necessary in our response framework.

Keywords: response theory; Fokker–Planck equation; overdamped limits; underdamped Langevin dynamics



Academic Editor: Michel Planat

Received: 9 July 2025

Revised: 4 August 2025

Accepted: 7 August 2025

Published: 11 August 2025

Citation: Colangeli, M.; Rondoni, L.; Vozza, P. Dissipation Functions and Brownian Oscillators. *Symmetry* **2025**, *17*, 1297. <https://doi.org/10.3390/sym17081297>

Copyright: © 2025 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Since the seminal contributions of Kubo [1,2], linear response theory has become a foundational framework in nonequilibrium statistical mechanics. Initially formulated to address Hamiltonian perturbations in deterministic systems, see [3], subsequent developments have significantly expanded its scope to incorporate nonlinear corrections, cf., e.g., [4,5], as well as non-Hamiltonian dynamics and dissipative stochastic processes [6–18], see also [19–21] for a historical perspective and [22–30] for experimental validations. The search for non-perturbative approaches to response theory, such as the method based on the transient time correlation function developed in [31], gained renewed momentum with the introduction of the *dissipation function*, defined in [32,33] as the energy dissipation rate that satisfies the fluctuation relation. This concept enabled the unification, within a consistent theoretical framework, of the diverse methodologies employed in nonequilibrium molecular dynamics. The Fluctuation Theorem, systematized in [34], provides a rigorous probabilistic description of entropy production in small, driven systems and applies to time-reversible dynamics subjected to constant external forces. Extensions to periodically driven systems were subsequently developed in [35–37]. The mathematical framework of response theory based on the dissipation function was further formalized in [38] and later extended to quantum dynamics [39], as well as to systems exhibiting collective behavior

and phase transitions, such as Kuramoto-type models [40], where standard linear response theory typically breaks down.

In this work, we extend this formalism to stochastic diffusion processes with additive noise. We demonstrate that many structural features originally derived for deterministic dynamics can be systematically transferred and adapted to stochastic systems governed by Fokker–Planck equations, see [41,42] for a general review.

We further derive explicit response formulae for both overdamped and underdamped Langevin dynamics. Importantly, these formulae are *exact*, in the sense that they are not limited to the linear response regime and remain valid for finite perturbation amplitudes.

Focusing on the analytically tractable case of harmonic Brownian oscillators, we illustrate the applicability of the considered response formalism, which may prove particularly useful in handling many-particle systems characterized by different time scales. We also examine the commutativity of two operations: the derivation of response expressions and the procedure leading to the overdamped limit, which corresponds to a perfect separation of time scales between fast and slow variables. This commutativity (or its absence) for selected observables reflects a structural symmetry of the response formalism (or a violation thereof) and may carry some practical implications. In particular, for systems with complex particle dynamics, it may be more tractable to probe the regime of perfect time scale separation at the level of observable averages, rather than by resolving the full stochastic dynamics of individual constituents. We also point out that the time symmetry of the microscopic dynamics, based on the existence of an involution in phase space [43], is not strictly required in the deterministic derivation of the response formalism outlined in [38]. Similarly, when adapting such a response formalism to Markovian dynamics, no explicit use of detailed balance is made, as illustrated, e.g., in Refs. [44–46].

This paper is organized as follows. In Section 2, we introduce the response theory based on the dissipation function. Section 3 presents the formal derivation of the overdamped limit of underdamped Langevin dynamics. In Section 4, we specialize to the harmonic case and derive exact response formulae for the overdamped dynamics of Brownian oscillators subjected to a perturbation that simultaneously alters the frequency of the external field and the temperature of the surrounding thermal bath. Section 5 addresses the underdamped case and analyzes the limiting behavior of the response in the regime of infinite time scale separation. Conclusions are drawn in Section 6, while technical details are provided in the Appendices A and B.

2. Dissipation Functions in the Fokker–Planck Equation

In this section, we analyze the role of the dissipation function in the evolution of probability densities governed by the Fokker–Planck equation. We begin with the stochastic differential equation (SDE) for a variable $x \in \mathbb{R}^d$ written in Itô's form:

$$dx(t) = A(x) dt + \sigma(x) dW(t), \quad x(0) = x_0, \quad (1)$$

where $A : \mathbb{R}^d \rightarrow \mathbb{R}^d$, $\sigma : \mathbb{R}^d \rightarrow \mathbb{R}^{d \times m}$ and $W(t)$ is a standard Wiener process in \mathbb{R}^m . The generator L of the process acts on state functions $O : \mathbb{R}^d \rightarrow \mathbb{R}$ as follows:

$$L O(x) = A(x) \cdot \nabla O(x) + D(x) : (\nabla \nabla O(x)), \quad (2)$$

where $D(x) = \frac{1}{2} \sigma(x) \sigma(x)^T$ is the diffusion matrix and $\nabla \nabla O$ denotes the Hessian matrix of O .

The L^2 -adjoint of the generator L , known as the *Fokker–Planck operator*, acts on probability densities $f(x, t)$ according to

$$L^* f(x, t) = \nabla \cdot [-A(x)p(x, t) + \nabla \cdot (D(x)f(x, t))]. \quad (3)$$

The corresponding forward Kolmogorov equation (or Fokker–Planck equation) for $f(x, t)$ reads as follows:

$$\begin{aligned} \partial_t f(x, t) &= L^* f(x, t), \quad \text{in } \mathbb{R}^d \times (0, \infty), \\ f(x, 0) &= f_0(x), \quad \text{on } \mathbb{R}^d \times \{t = 0\}. \end{aligned} \quad (4)$$

Using the shorthand notation $f_t(x) = f(x, t)$, Equation (4) can be rewritten as

$$\partial_t f_t(x) = \Omega^{f_t}(x) f_t(x), \quad (5)$$

where the state function $\Omega^{f_t}(x) := (L^* f_t(x))/f_t(x)$, called the dissipation function (note that we purposely adopt, here, the same terminology used in the deterministic setting [38]), is given explicitly by

$$\begin{aligned} \Omega^{f_t}(x) &= -\nabla \cdot A(x) - A(x) \cdot \nabla \log f_t(x) + \nabla \nabla : D(x) \\ &\quad + 2(\nabla \cdot D(x)) \cdot \nabla \log f_t(x) \\ &\quad + D(x) : [\nabla \nabla \log f_t(x) + \nabla \log f_t(x) \otimes \nabla \log f_t(x)], \end{aligned} \quad (6)$$

see Appendix A for details. Finally, we define the expectation of the state function $O(x)$ at time $s \geq 0$ as follows:

$$\langle O \rangle_s = \int_{\mathbb{R}^d} O(x) f_s(x) dx. \quad (7)$$

Exploiting the formal solution of Equation (4), we find

$$\langle O \rangle_s = \int O(x) (e^{sL^*} p_0(x)) dx = \int (e^{sL} O(x)) p_0(x) dx.$$

The time derivative of $\langle O \rangle_s$ reads

$$\frac{d}{ds} \langle O \rangle_s = \int \left(\frac{d}{ds} e^{sL} O(x) \right) f_0(x) dx = \int (L e^{sL} O(x)) f_0(x) dx,$$

which, using the adjoint operator, becomes the following:

$$\frac{d}{ds} \langle O \rangle_s = \int (e^{sL} O(x)) (L^* f_0(x)) dx.$$

Finally, from Equation (5), we find

$$\frac{d}{ds} \langle O \rangle_s = \int (e^{sL} O(x)) \Omega^{f_0}(x) f_0(x) dx = \langle O(x_s) \Omega^{f_0}(x_0) \rangle_0. \quad (8)$$

By integrating both sides of (8) from 0 to t , we finally obtain the exact response formula:

$$\langle O \rangle_t - \langle O \rangle_0 = \int_0^t \langle O(x_s) \Omega^{f_0}(x_0) \rangle_0 ds. \quad (9)$$

We note that the average appearing under the integral in Equation (9) is taken with respect to the initial density $f_0(x)$. To establish a connection with response theory, it is helpful to further characterize $f_0(x)$ introducing an alternative process, called the reference dynamics and defined for times $t \in (-\infty, 0)$, which admits $f_0(x)$ as its invariant probability density. The dynamics governed by Equation (1) can be interpreted as the result of applying a

perturbation to the reference dynamics at time $t = 0$ and therefore will be referred to as the perturbed dynamics.

We also observe that if in Equation (9) one sets $O(x) = 1$ for all $x \in \mathbb{R}^d$, then the normalization of the probability density $f_s(x)$, for $s \geq 0$, implies $\langle \Omega^{f_0} \rangle_0 = 0$. We finally remark that, unlike classical linear response formulae for Markovian diffusions, as discussed, for instance, in [47–49], formula (9) remains valid even in the presence of perturbations that drive the process arbitrarily far from the reference dynamics.

3. From the Underdamped to the Overdamped Langevin Dynamics

We let $x = (q, p)^T \in \mathbb{R}^2$ and consider the underdamped Langevin equation describing a Brownian particle subjected to an external potential $U(q)$, see [47,48]:

$$dq(t) = \frac{p}{m} dt, \tag{10a}$$

$$dp(t) = -\nabla_q U(q) dt - \gamma p dt + \sqrt{2\gamma m \beta^{-1}} dW(t). \tag{10b}$$

We briefly outline the formal steps leading to the overdamped limit of Equation (10), following the standard derivation described in [48,50]. We introduce $\epsilon > 0$, and in Equation (10) we rescale $\gamma \mapsto \gamma/\epsilon$ and $t \mapsto t/\epsilon$, which yields

$$dq^\epsilon = \frac{p^\epsilon}{\epsilon m} dt, \tag{11a}$$

$$dp^\epsilon = -\frac{1}{\epsilon} \nabla_q U(q^\epsilon) dt - \frac{\gamma}{\epsilon^2} p^\epsilon dt + \frac{1}{\epsilon} \sqrt{2\gamma m \beta^{-1}} dW(t), \tag{11b}$$

where we have exploited the scaling property of the Wiener process $W(t/\epsilon) = 1/\sqrt{\epsilon} W(t)$. From Equations (11a) and (11b), we obtain

$$dq^\epsilon = \frac{p^\epsilon}{\epsilon m} dt = -\frac{\epsilon}{\gamma m} dp^\epsilon - \frac{1}{\gamma m} \nabla_q U(q^\epsilon) dt + \sqrt{\frac{2}{m\beta\gamma}} dW(t).$$

Taking the limit $\epsilon \rightarrow 0$, we arrive at the overdamped Langevin dynamics [48]

$$dq = -\frac{1}{\gamma m} \nabla_q U(q) dt + \sqrt{\frac{2}{m\beta\gamma}} dW(t). \tag{12}$$

In the sequel we will focus on Brownian oscillators, namely Brownian particles bound by a harmonic potential $U(q) = \frac{1}{2} m \omega^2 q^2$, and evaluate Equation (9) for both the overdamped dynamics in Equation (12) (in Section 4) and its underdamped counterpart expressed by Equation (10) (in Section 5).

4. Exact Response Theory for the Overdamped Langevin Dynamics

The application of the response theory introduced in Section 2 to Equation (12) begins with the definition of the reference dynamics:

$$dq(t) = -\frac{\omega_0^2 q}{\gamma} dt + \sqrt{\frac{2}{m\beta_0\gamma}} dW(t), \quad t \leq 0, \tag{13}$$

which switches, at time $t = 0^+$, into the perturbed one, defined as follows:

$$dq(t) = -\frac{\omega^2 q}{\gamma} dt + \sqrt{\frac{2}{m\beta\gamma}} dW(t), \quad t > 0. \tag{14}$$

Note that both parameters $\omega_0 \rightarrow \omega$ and $\beta_0 \rightarrow \beta$ have changed as a result of the perturbation. The Fokker–Planck equation corresponding to Equation (14), known as the Smoluchowski equation [49], reads

$$\frac{\partial f_t(q)}{\partial t} = \frac{\partial}{\partial q} \left[\frac{\omega^2}{\gamma} q f_t(q) + \frac{1}{m\beta\gamma} \frac{\partial}{\partial q} f_t(q) \right], \quad (15)$$

and is supplied with the initial datum

$$f_0(q) = \sqrt{\frac{m\beta_0\omega_0^2}{2\pi}} \exp\left(-\frac{1}{2}m\beta_0\omega_0^2q^2\right). \quad (16)$$

Equation (15) can be cast in the form

$$\frac{\partial f_t(q)}{\partial t} = \Omega_{\omega,\beta}^{f_t} f_t, \quad (17)$$

where the subscripts in $\Omega_{\omega,\beta}^{f_t}$ highlight the two parameters that characterize the perturbation. A straightforward computation (see Equation (A5) in Appendix A) leads to the following expression of the dissipation function $\Omega_{\omega,\beta}^{f_0}$ characterizing the overdamped dynamics (14):

$$\Omega_{\omega,\beta}^{f_0}(q) = A_0 + B_0q^2, \quad (18)$$

where $A_0 = (\beta\omega^2 - \beta_0\omega_0^2)/(\beta\gamma)$ and $B_0 = -A_0/\langle q^2 \rangle_0$, with $\langle q^2 \rangle_0 = (\beta_0m\omega_0^2)^{-1}$. Formula (18) reveals that the initial distribution (16) is effectively invariant for the reference dynamics (13), because the following holds:

$$\Omega_{\omega_0,\beta_0}^{f_0}(q) = 0. \quad (19)$$

Moreover, we also point out that $\Omega_{\omega,\beta}^{f_0}$ vanishes, more generally, if the condition $\beta\omega^2 = \beta_0\omega_0^2$ is satisfied. This follows from the fact that the invariant densities of the reference and the perturbed dynamics are both Gaussian distributions:

$$f_0(q) = \mathcal{N}\left(0, \frac{1}{\beta_0m\omega_0^2}\right), \quad f_\infty(q) = \mathcal{N}\left(0, \frac{1}{\beta m\omega^2}\right).$$

Under the condition $\beta\omega^2 = \beta_0\omega_0^2$, the variances of these distributions coincide (and so do the respective means, which both vanish), implying $f_0(q) \equiv f_\infty(q)$. For the Brownian oscillator model described by Equation (14), the expression of the conditional probability density $p(r, t|q, 0)$ can be made explicit:

$$p(r, t|q, 0) = \sqrt{\frac{\beta m\omega^2}{2\pi\left(1 - e^{-\frac{2\omega^2}{\gamma}t}\right)}} \exp\left[-\frac{\beta m\omega^2\left(r - q e^{-\frac{\omega^2}{\gamma}t}\right)^2}{2\left(1 - e^{-\frac{2\omega^2}{\gamma}t}\right)}\right]. \quad (20)$$

We compute, first, the exact response formula for the observable $O(q) \equiv q^2$. In this case Equation (9) reads

$$\langle q^2 \rangle_t = \langle q^2 \rangle_0 + \int_0^t \langle q_s^2 \Omega_{\omega,\beta}^{f_0}(q_0) \rangle_0 ds. \quad (21)$$

From Equation (20), we obtain the expression

$$\langle q^2 \rangle_t = \frac{\beta\omega^2 e^{-2\kappa t} + (1 - e^{-2\kappa t})\beta_0\omega_0^2}{m\beta_0\beta\omega_0^2\omega^2}, \quad (22)$$

where $\kappa = \omega^2/\gamma$ denotes a characteristic relaxation frequency. We point out that formula (22) possesses the proper asymptotic behavior: it reduces to $\langle q^2 \rangle_0$ at $t = 0$ while it recovers the expression $\langle q^2 \rangle_\infty = (\beta m \omega^2)^{-1}$ in the long-time limit. We turn, then, to the analysis of the response formula for the state function $O(q) \equiv \Omega_{\omega,\beta}^{f_0}(q)$. In this case formula (9) reads

$$\langle \Omega_{\omega,\beta}^{f_0} \rangle_t = \langle \Omega_{\omega,\beta}^{f_0} \rangle_0 + \int_0^t \langle \Omega_{\omega,\beta}^{f_0}(q_s) \Omega_{\omega,\beta}^{f_0}(q_0) \rangle_0 ds. \quad (23)$$

We first verify that it correctly satisfies

$$\langle \Omega_{\omega,\beta}^{f_0} \rangle_0 = A_0 + B_0 \langle q^2 \rangle_0 = 0, \quad (24)$$

whereas

$$\begin{aligned} \int_0^t \langle \Omega_{\omega,\beta}^{f_0}(q_s) \Omega_{\omega,\beta}^{f_0}(q_0) \rangle_0 ds &= B_0 \int_0^s \langle q_s^2, \Omega_{\omega,\beta}^{f_0}(q_0) \rangle_0 ds \\ &= B_0 (1 - e^{-2\kappa t}) (\langle q^2 \rangle_\infty - \langle q^2 \rangle_0) \\ &= \frac{(\beta \omega^2 - \beta_0 \omega_0^2)^2}{\beta^2 \gamma \omega^2} (1 - e^{-2\kappa t}). \end{aligned} \quad (25)$$

5. Exact Response Theory for the Underdamped Langevin Dynamics

We now apply the formalism of Section 2 in the setup of the underdamped dynamics expressed by Equation (10). We start, as in Section 4, with the definition of the reference dynamics for $t \leq 0$, which takes the form

$$dq(t) = \frac{p}{m} dt, \quad (26a)$$

$$dp(t) = -m\omega_0^2 q dt - \gamma p dt + \sqrt{2\gamma m \beta_0^{-1}} dW(t). \quad (26b)$$

At time $t = 0^+$ the joint perturbation $\omega_0 \rightarrow \omega$ and $\beta_0 \rightarrow \beta$ makes the reference dynamics turn into the perturbed one, defined for $t > 0$ as follows:

$$dq(t) = \frac{p}{m} dt, \quad (27a)$$

$$dp(t) = -m\omega^2 q dt - \gamma p dt + \sqrt{2\gamma m \beta^{-1}} dW(t). \quad (27b)$$

The Fokker–Planck equation describing, for $t > 0$, the evolution of the probability density $f_t(q, p)$ (an instance of the Kramers equation [49]) reads

$$\frac{\partial f_t}{\partial t} = -\frac{p}{m} \frac{\partial f_t}{\partial q} + m\omega_0^2 q \frac{\partial f_t}{\partial p} + \gamma f_t + \gamma p \frac{\partial f_t}{\partial p} + \frac{\gamma m}{\beta} \frac{\partial^2 f_t}{\partial p^2}, \quad (28)$$

and is supplied with the initial datum

$$f_0(q, p) = \frac{\beta_0 \omega_0}{2\pi} e^{-\frac{\beta_0}{2} \left(m\omega_0^2 q^2 + \frac{p^2}{m} \right)}. \quad (29)$$

The dissipation function $\Omega_{\omega,\beta}^{f_0}$ derived from (15) corresponds to the state function

$$\Omega_{\omega,\beta}^{f_0}(q, p) = A_1 p^2 + B_1 q p + C_1, \quad (30)$$

where $A_1 = -\frac{\gamma\beta_0}{m}\left(1 - \frac{\beta_0}{\beta}\right)$, $B_1 = \beta_0(\omega_0^2 - \omega^2)$ and $C_1 = -A_1\langle p^2 \rangle_0$, with $\langle p^2 \rangle_0 = m/\beta_0$. We note that $f_0(q, p)$ in (29) is invariant for the reference dynamics (26), for $\Omega_{\omega_0, \beta_0}^{f_0} = 0$ holds. Furthermore, letting $x = (q, p)^T \in \mathbb{R}^2$, Equation (28) can be rewritten in the form

$$\frac{\partial f_t}{\partial t} = \sum_{i,j=1}^2 \left[-A_{i,j} \frac{\partial}{\partial x_i} (x_j f_t) + D_{i,j} \frac{\partial^2 f_t}{\partial x_i \partial x_j} \right], \tag{31}$$

where the matrices A and D have the following form:

$$A = \begin{pmatrix} 0 & 1/m \\ -m\omega^2 & -\gamma \end{pmatrix}, \quad D = \begin{pmatrix} 0 & 0 \\ 0 & \gamma/\beta \end{pmatrix}. \tag{32}$$

In the large friction regime $\gamma > 2\omega$ that is considered here in the setup of the underdamped Langevin model, the two eigenvalues of the matrix A are real, $-\lambda_{\pm} < 0$, with $\lambda_{\pm} = (\gamma \pm \sqrt{\gamma^2 - 4\omega^2})/2$. The time scale separation, in this model, is thus quantified by $\Delta = \lambda_+ - \lambda_- = \sqrt{\gamma^2 - 4\omega^2}$.

Proceeding as in Section 4, let us consider, first, the exact response formula for the observable $O(q, p) \equiv q^2$.

The evaluation of time correlation functions for the underdamped dynamics (27) $\langle O(x_s) \Omega_{\omega\beta}^{f_0}(x_0) \rangle_0$ can be carried out by knowledge of the conditional probability density $p(y, t|x, 0)$, which in this case reads

$$p(y, t|x, 0) = \frac{1}{2\pi\sqrt{\det \Sigma(t)}} \exp\left(-\frac{1}{2}(y - e^{At}x)\Sigma^{-1}(t)(y - e^{At}x)\right), \tag{33}$$

where

$$\Sigma(t) = \int_0^t e^{As} D e^{A^T s} ds$$

is the covariance matrix. We point out that the stationary covariance matrix $\bar{\Sigma} := \lim_{t \rightarrow \infty} \Sigma(t)$ satisfies the Lyapunov equation.

$$A\bar{\Sigma} + \bar{\Sigma}A^T = -D,$$

which is an instance of the fluctuation–dissipation relation, see, e.g., [47,48,50]. Using Equation (A6e) in Appendix B, we find that, for the state function $O(q, p) \equiv q^2$, formula (9) evaluates to

$$\begin{aligned} \langle q^2 \rangle_t &= \frac{1}{\beta_0 m \omega_0^2} + \frac{e^{-2t(\lambda_- + \lambda_+)}}{m\beta\beta_0\lambda_-(\lambda_- - \lambda_+)^2\lambda_+(\lambda_- + \lambda_+)\omega_0^2} \times \\ &\quad \left[e^{2t\lambda_-} \lambda_-(\lambda_- + \lambda_+) \left(\beta\lambda_- \omega^2 - (\beta_0\gamma + \beta(-\gamma + \lambda_-))\omega_0^2 \right) \right. \\ &\quad + e^{2t\lambda_+} \lambda_+(\lambda_- + \lambda_+) \left(\beta\lambda_+ \omega^2 - (\beta_0\gamma + \beta(-\gamma + \lambda_+))\omega_0^2 \right) \\ &\quad - 2e^{t(\lambda_- + \lambda_+)} \lambda_- \lambda_+ \left(\beta(\lambda_- + \lambda_+) \omega^2 - (2\beta_0\gamma + \beta(-2\gamma + \lambda_- + \lambda_+))\omega_0^2 \right) \\ &\quad + e^{2t(\lambda_- + \lambda_+)} (\lambda_- - \lambda_+)^2 \times \\ &\quad \left. \left(-\beta(\lambda_- + \lambda_+) \omega^2 + (\beta_0\gamma + \beta(-\gamma + \lambda_- + \lambda_+))\omega_0^2 \right) \right]. \tag{34} \end{aligned}$$

More insight into formula (34) is gained by inspecting the large- γ regime. To this aim we adopt in Equation (34) the scaling previously introduced in Section 2, i.e., $\gamma \rightarrow \gamma/\epsilon$, $t \rightarrow t/\epsilon$. The following asymptotics holds:

$$\lambda_+ = \frac{\gamma}{\epsilon} + O(\epsilon) \quad , \quad \lambda_- = \frac{\omega^2}{\gamma} \epsilon + o(\epsilon). \quad (35)$$

Note that the time scale separation $\Delta \sim \epsilon^{-1}$ diverges as $\epsilon \rightarrow 0$, which corresponds to the regime of perfect time scale separation. Inserting the expressions (35) in (34) and then taking the overdamped limit $\epsilon \rightarrow 0$, one obtains

$$\langle q^2 \rangle_t = \frac{\beta \omega^2 e^{-2\frac{\omega^2}{\gamma}t} + \left(1 - e^{-2\frac{\omega^2}{\gamma}t}\right) \beta_0 \omega_0^2}{m \beta_0 \beta \omega_0^2 \omega^2}, \quad (36)$$

which recovers formula (22). This result highlights a key feature of the response theory developed in Section 2: for certain observables $O(q, p)$, the two operations of taking the limit $\epsilon \rightarrow 0$ and deriving the response formula (9) commute.

We now turn our attention to the state function $O(q, p) = p^2$. In this context, the momentum $p(t)$ is regarded as a *fast* variable, while the rescaling $t \rightarrow t/\epsilon$ corresponds to observing the evolution of the system on a *slow* time scale [51]. On such a scale, as ϵ decreases, the quantity $\langle p^2 \rangle_t$ is expected to thermalize increasingly rapidly toward its equilibrium value m/β , as dictated by the equipartition theorem. This behavior is fully confirmed by our calculations, which, using Equation (A6d) in Appendix B, show that

$$\begin{aligned} \langle p^2 \rangle_t &= \frac{1}{\beta_0 m} \left(1 + \frac{e^{-2t(\lambda_- + \lambda_+)} (\beta - \beta_0) \gamma}{\beta (\lambda_- - \lambda_+)^2 (\lambda_- + \lambda_+)} \left[-e^{2t(\lambda_- + \lambda_+)} (\lambda_- - \lambda_+)^2 \right. \right. \\ &\quad \left. \left. - 4e^{t(\lambda_- + \lambda_+)} \lambda_- \lambda_+ + (\lambda_- + \lambda_+) (e^{2t\lambda_+} \lambda_- + e^{2t\lambda_-} \lambda_+) \right] \right. \\ &\quad \left. + \frac{e^{-2t(\lambda_- + \lambda_+)} (e^{t\lambda_-} - e^{t\lambda_+})^2 \omega^2 (\omega - \omega_0) (\omega + \omega_0)}{(\lambda_- - \lambda_+)^2 \omega_0^2} \right). \end{aligned} \quad (37)$$

At $t = 0$ formula (37) correctly reproduces $\langle p^2 \rangle_0 = m/\beta_0$. Letting $\gamma \rightarrow \gamma/\epsilon$, $t \rightarrow t/\epsilon$, using (35), and finally passing to the limit $\epsilon \rightarrow 0$, we find

$$\langle p^2 \rangle_t = \frac{m}{\beta} \quad , \quad t > 0, \quad (38)$$

which confirms that, after the rescaling, the quantity $\langle p^2 \rangle_t$ undergoes an instantaneous thermalization.

As a next step, we consider the state function $O(q, p) = qp$. Using Equation (A6c) in Appendix B, we find

$$\begin{aligned} \langle qp \rangle_t &= \frac{e^{-2t(\lambda_- + \lambda_+)}}{2\beta\beta_0\lambda_-(\lambda_- - \lambda_+)^2\lambda_+(\lambda_- + \lambda_+)\omega_0^2} \times \\ &\quad \left[-e^{2t(\lambda_- + \lambda_+)} \beta (\lambda_- - \lambda_+)^2 (\lambda_- \lambda_+ - \omega^2) (\omega^2 - \omega_0^2) \right. \\ &\quad \left. - e^{2t\lambda_+} \lambda_+ (\lambda_- + \lambda_+) [\beta \omega^2 (\lambda_- \lambda_+ + \omega^2)] \right. \\ &\quad \left. - (2\beta_0 \gamma \lambda_- + \beta (-2\gamma \lambda_- + \lambda_- \lambda_+ + \omega^2)) \omega_0^2 \right] \\ &\quad \left[-e^{2t\lambda_-} \lambda_- (\lambda_- + \lambda_+) [\beta \omega^2 (\lambda_- \lambda_+ + \omega^2)] \right. \\ &\quad \left. - (2\beta_0 \gamma \lambda_+ + \beta (-2\gamma \lambda_+ + \lambda_- \lambda_+ + \omega^2)) \omega_0^2 \right] \\ &\quad \left. + 2e^{t(\lambda_- + \lambda_+)} \lambda_- \lambda_+ [\beta \omega^2 (\lambda_-^2 + \lambda_+^2 + 2\omega^2)] \right. \\ &\quad \left. - (2\beta_0 \gamma (\lambda_- + \lambda_+) + \beta (\lambda_-^2 + \lambda_+^2 - 2\gamma (\lambda_- + \lambda_+) + 2\omega^2)) \omega_0^2 \right], \end{aligned} \quad (39)$$

which, after rescaling $\gamma \rightarrow \gamma/\epsilon$, $t \rightarrow t/\epsilon$ and passing to the limit $\epsilon \rightarrow 0$, yields

$$\langle qp \rangle_t = 0 \quad , \quad t > 0. \quad (40)$$

Finally, we consider the state function $O(q, p) \equiv \Omega_{\omega, \beta}^{f_0}(q, p)$. From Equation (30) we can write

$$\langle \Omega_{\omega, \beta}^{f_0} \rangle_t = A_1 \langle p^2 \rangle_t + B_1 \langle qp \rangle_t + C_1. \quad (41)$$

At $t = 0$ one correctly finds

$$\langle \Omega_{\omega, \beta}^{f_0} \rangle_t = A_1 \langle p^2 \rangle_0 + C_1 = 0.$$

Introducing the usual scaling $\gamma \rightarrow \gamma/\epsilon$, $t \rightarrow t/\epsilon$ in the limit $\epsilon \rightarrow 0$, we obtain

$$\begin{aligned} \langle \Omega_{\omega, \beta}^{f_0} \rangle_t &= A_1 \langle p^2 \rangle_t + C_1 \quad , \quad t > 0 \\ &= \frac{\gamma}{\beta^2} (\beta - \beta_0)^2. \end{aligned} \quad (42)$$

We observe that the quantity $\langle \Omega_{\omega, \beta}^{f_0} \rangle_t$ undergoes an instantaneous thermalization in the appropriate scaling limit: it exhibits a discontinuous jump from the value 0 at $t = 0$ to the final value expressed by Equation (42).

We further note that Equation (42) does not recover the expression in Equation (25). This discrepancy arises because the observables defined in Equations (18) and (30), although both referred to as dissipation functions, correspond to distinct state functions.

6. Conclusions

In this work, we have extended the response theory based on dissipation functions, originally developed in the context of deterministic dissipative dynamical systems [38], to Markovian diffusion processes governed by a Fokker–Planck equation. A key feature of our derivation is twofold: (i) the response formulae are conveniently expressed in terms of the initial probability density [38] and (ii) these formulae remain valid for perturbations of arbitrary magnitude and are therefore not restricted to the conventional linear response regime. We also note that in our derivation the time reversal symmetry often assumed in statistical mechanics has not been used, although it can of course be present. This is of interest in the investigation of the effects of symmetries in systems of physical interest. Focusing on the analytically tractable case of particles confined in a harmonic potential and subjected to additive noise, we have demonstrated a notable property of the formalism: for specific observables, the operations of deriving the response formulae and taking the limit $\epsilon \rightarrow 0$, known as the overdamped limit, commute. Moreover, the theory correctly captures the average behavior of certain state functions, such as the dissipation function associated with the underdamped dynamics, which exhibits instantaneous relaxation to its equilibrium value in the overdamped limit, see Equation (42). While this behavior is consistent with model reduction procedures such as adiabatic elimination, which rely on a large separation of time scales [42,52], it does not reproduce the behavior of the average dissipation function obtained for the overdamped dynamics, see Equation (25). This discrepancy arises from the fact that the two dissipation functions depend on different sets of variables, namely fast variables in the underdamped case and slow variables in the overdamped dynamics, see Equations (18) and (30), respectively. As a result, their averages exhibit distinct behavior in the overdamped limit.

Although such discrepancies are often encountered in the study of anomalous transport phenomena [53], they can more fundamentally be traced back to the distinct time scales governing the decay of correlation functions, as explicitly derived for the Brownian oscillator model in Sections 4 and 5.

Looking ahead, the framework developed in this work sets the stage for applying response theory to high-dimensional Markovian diffusion processes, which may exhibit a cascade of different time scales. In these more complex models, for which analytical solutions may not be available, our formalism can still be exploited via Monte Carlo simulations. Furthermore, model reduction techniques can assist in identifying a relevant subset of resolved variables, typically those evolving on slow time scales and governing the macroscopic behavior of the system.

Author Contributions: Conceptualization, M.C., L.R. and P.V.; methodology, M.C., L.R. and P.V.; software, P.V.; validation, M.C., L.R. and P.V.; formal analysis, M.C., L.R. and P.V.; investigation, M.C., L.R. and P.V.; data curation, P.V.; writing—original draft preparation, M.C.; writing—review and editing, L.R. and P.V.; visualization, P.V.; supervision, M.C. and L.R. All authors have read and agreed to the published version of the manuscript.

Funding: This work was carried out under the auspices of the Italian National Group of Mathematical Physics. The research of MC has been developed in the framework of the research project National Centre for HPC, Big Data and Quantum Computing—PNRR Project, funded by the European Union—NextGenerationEU.

Data Availability Statement: No new data were created or analyzed in this study. Data sharing is not applicable to this article.

Conflicts of Interest: The authors declare no conflicts of interest.

Appendix A. Derivation of the Dissipation Function for Fokker–Planck Dynamics

We sketch the derivation of the dissipation function $\Omega^{f_t}(x) : \mathbb{R}^d \rightarrow \mathbb{R}$, defined as follows:

$$\Omega^{f_t}(x) = \frac{L^* f_t(x)}{f_t(x)},$$

expressed in terms of the Fokker–Planck operator L^* in \mathbb{R}^d , i.e.,

$$L^* f_t(x) = \nabla \cdot [-A(x)f_t(x) + \nabla \cdot (D(x)f_t(x))], \quad (\text{A1})$$

cf. Equation (5). Using the identities

$$\begin{aligned} \nabla \cdot (A(x)f_t(x)) &= (\nabla \cdot A(x))f_t(x) + A(x) \cdot \nabla f_t(x) \\ \nabla \cdot (\nabla \cdot (D(x)f_t(x))) &= \nabla \nabla : (D(x)f_t(x)), \end{aligned}$$

we obtain

$$L^* f = -(\nabla \cdot A(x))f_t(x) - A(x) \cdot \nabla f_t(x) + \nabla \nabla : (D(x)f_t(x)),$$

which yields

$$\Omega^{f_t}(x) = -\nabla \cdot A(x) - A(x) \cdot \nabla \log f_t(x) + \frac{1}{f_t(x)} \nabla \nabla : (D(x)f_t(x)).$$

Then, expanding the last term

$$\frac{1}{f_t(x)} \nabla \nabla : (D(x)f_t(x)) = \nabla \nabla : D(x) + 2 \frac{(\nabla \cdot D(x)) \cdot \nabla f_t(x)}{f_t(x)} + \frac{D(x) : \nabla \nabla f_t(x)}{f_t(x)}$$

and using the identity

$$\frac{\nabla f_t(x)}{f_t(x)} = \nabla \log f_t(x), \quad \frac{\nabla \nabla f_t(x)}{f_t(x)} = \nabla \nabla \log f_t(x) + \nabla \log f_t(x) \otimes \nabla \log f_t(x),$$

we obtain

$$\begin{aligned} \Omega^{f_t}(x) &= -\nabla \cdot A(x) - A(x) \cdot \nabla \log f_t(x) + \nabla \nabla : D(x) \\ &\quad + 2(\nabla \cdot D(x)) \cdot \nabla \log f_t(x) \\ &\quad + D(x) : [\nabla \nabla \log f_t(x) + \nabla \log f_t(x) \otimes \nabla \log f_t(x)], \end{aligned} \tag{A2}$$

which recovers Equation (6).

In dimension $d = 1$ Equation (A2) reduces to

$$\Omega^{f_t}(x) = -A'(x) - A(x) \frac{f_t'(x)}{f_t(x)} + D''(x) + 2D'(x) \frac{f_t'(x)}{f_t(x)} + D(x) \frac{f_t''(x)}{f_t(x)}. \tag{A3}$$

Furthermore, in the presence of constant $A(x) = A$ and $D(x) = D$, Equation (A2) yields

$$\Omega^{p_t}(x) = -A \cdot \nabla \log f_t(x) + D : \nabla \nabla \log f_t(x) + D : (\nabla \log f_t(x) \otimes \nabla \log f_t(x)). \tag{A4}$$

which admits an even further simplification for $d = 1$:

$$\Omega^{f_t}(x) = -A \frac{f_t'(x)}{f_t(x)} + D \frac{f_t''(x)}{f_t(x)}. \tag{A5}$$

Appendix B. Conditional Averages for the Underdamped Dynamics

Using the setup introduced in Section 5, let $O : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a generic state function and define

$$\bar{O}(x, t) = \int O(y) p(y, t | x, 0) dy$$

as its conditional expectation, computed with respect to the conditional probability density $p(y, t | x, 0)$, as given in Equation (33). From the analysis of Chandrasekhar [54], in the regime $\gamma > 2\omega$ one obtains the following results:

$$\bar{q}_t = e^{-\frac{\gamma t}{2}} \left(\cosh\left(\frac{\Delta t}{2}\right) + \frac{\gamma}{\Delta} \sinh\left(\frac{\Delta t}{2}\right) \right) q + \frac{2}{\Delta} e^{-\frac{\gamma t}{2}} \sinh\left(\frac{\Delta t}{2}\right) \frac{p}{m} \tag{A6a}$$

$$\bar{p}_t = -\frac{2m\omega^2}{\Delta} e^{-\frac{\gamma t}{2}} \sinh\left(\frac{\Delta t}{2}\right) q + e^{-\frac{\gamma t}{2}} \left(\cosh\left(\frac{\Delta t}{2}\right) - \frac{\gamma}{\Delta} \sinh\left(\frac{\Delta t}{2}\right) \right) p \tag{A6b}$$

$$\bar{q}_t \bar{p}_t = \bar{q}_t \bar{p}_t + \frac{4}{\beta} \frac{\gamma}{\Delta^2} e^{-\gamma t} \sinh^2\left(\frac{\Delta t}{2}\right) \tag{A6c}$$

$$\bar{p}_t^2 = \bar{p}_t^2 + \frac{m}{\beta} \left(1 - e^{-\gamma t} \left(2\left(\frac{\gamma}{\Delta}\right)^2 \sinh^2\left(\frac{\Delta t}{2}\right) - \frac{\gamma}{\Delta} \sinh(\Delta t) + 1 \right) \right) \tag{A6d}$$

$$\bar{q}_t^2 = \bar{q}_t^2 + \frac{1}{m\beta\omega^2} \left(1 - e^{-\gamma t} \left(2\left(\frac{\gamma}{\Delta}\right)^2 \sinh^2\left(\frac{\Delta t}{2}\right) + \frac{\gamma}{\Delta} \sinh(\Delta t) + 1 \right) \right), \tag{A6e}$$

with $\Delta = \lambda_+ - \lambda_-$.

References

1. Kubo, R. The Fluctuation-Dissipation Theorem. *Rep. Prog. Phys.* **1966**, *29*, 255–284. [\[CrossRef\]](#)
2. Kubo, R.; Toda, M.; Hashitsume, N. *Statistical Physics II. Nonequilibrium Statistical Mechanics*; Springer: Berlin, Germany, 1985.
3. Zwanzig, R. *Nonequilibrium Statistical Mechanics*; Oxford University Press: Oxford, UK, 2001.

4. Bochkov, G.N.; Kuzovlev, Y.E. Nonlinear fluctuation-dissipation relations and stochastic models in nonequilibrium thermodynamics: I. Generalized fluctuation-dissipation theorem. *Phys. A* **1981**, *106*, 443–479. [[CrossRef](#)]
5. Lucarini, V.; Colangeli, M. Beyond the linear fluctuation-dissipation theorem: The role of causality. *J. Stat. Mech. Theory Exp.* **2012**, *2012*, P05013. [[CrossRef](#)]
6. Ruelle, D. General linear response formula in statistical mechanics, and the fluctuation-dissipation theorem far from equilibrium. *Phys. Lett. A* **1998**, *245*, 220–224. [[CrossRef](#)]
7. Ruelle, D. Statistical mechanics: Rigorous results and applications to dynamical systems far from equilibrium. *Phys. Rep.* **1999**, *303*, 259–281. [[CrossRef](#)]
8. Marconi, U.M.B.; Puglisi, A.; Rondoni, L.; Vulpiani, A. Fluctuation–dissipation: Response theory in statistical physics. *Phys. Rep.* **2008**, *461*, 111–195. [[CrossRef](#)]
9. Prost, J.; Joanny, J.F.; Parrondo, J.M.R. Generalized fluctuation-dissipation theorem for steady-state systems. *Phys. Rev. Lett.* **2009**, *103*, 090601. [[CrossRef](#)] [[PubMed](#)]
10. Baiesi, M.; Maes, C.; Wynants, B. Fluctuations and response of nonequilibrium states. *Phys. Rev. Lett.* **2009**, *103*, 010602. [[CrossRef](#)] [[PubMed](#)]
11. Baiesi, M.; Maes, C.; Wynants, B. Nonequilibrium Linear Response for Markov Dynamics, I: Jump Processes and Overdamped Diffusions. *J. Stat. Phys.* **2009**, *137*, 1094–1116. [[CrossRef](#)]
12. Baiesi, M.; Boksenbojm, E.; Maes, C.; Wynants, B. Nonequilibrium Linear Response for Markov Dynamics, II: Inertial Dynamics. *J. Stat. Phys.* **2010**, *139*, 492–505. [[CrossRef](#)]
13. Seifert, U.; Speck, T. Fluctuation-dissipation theorem in nonequilibrium steady states. *Europhys. Lett.* **2010**, *89*, 10007. [[CrossRef](#)]
14. Colangeli, M.; Maes, C.; Wynants, B. A meaningful expansion around detailed balance. *J. Phys. A Math. Theor.* **2011**, *44*, 095001. [[CrossRef](#)]
15. Colangeli, M.; Rondoni, L.; Vulpiani, A. Fluctuation-dissipation relation for chaotic non-Hamiltonian systems. *J. Stat. Mech. Theory Exp.* **2012**, *2012*, L04002. [[CrossRef](#)]
16. Maes, C. Response Theory: A Trajectory-Based Approach. *Front. Phys.* **2020**, *8*, 229. [[CrossRef](#)]
17. Maes, C. Frenesy: Time-symmetric dynamical activity in nonequilibria. *Phys. Rep.* **2020**, *850*, 1–33. [[CrossRef](#)]
18. Baldovin, M.; Caprini, L.; Puglisi, A.; Sarracino, A.; Vulpiani, A. The Many Faces of Fluctuation–Dissipation Relations Out of Equilibrium. In *Nonequilibrium Thermodynamics and Fluctuation Kinetics*; Springer: Berlin/Heidelberg, Germany, 2022; pp. 47–79. [[CrossRef](#)]
19. Nyquist, H. Thermal Agitation of Electric Charge in Conductors. *Phys. Rev.* **1928**, *32*, 110–113. [[CrossRef](#)]
20. Johnson, J.B. Thermal Agitation of Electricity in Conductors. *Phys. Rev.* **1928**, *32*, 97–109. [[CrossRef](#)]
21. Callen, H.B.; Welton, T.A. Irreversibility and Generalized Noise. *Phys. Rev.* **1951**, *83*, 34–40. [[CrossRef](#)]
22. Bellon, L.; Ciliberto, S.; Laroche, C. Violation of the fluctuation-dissipation relation during the formation of a colloidal glass. *Europhys. Lett.* **2001**, *53*, 511–517. [[CrossRef](#)]
23. Wang, G.M.; Sevick, E.M.; Mittag, E.; Searles, D.J.; Evans, D.J. Experimental demonstration of violations of the second law of thermodynamics for small systems and short time scales. *Phys. Rev. Lett.* **2002**, *89*, 050601. [[CrossRef](#)]
24. Ojha, R.P.; Lemieux, P.A.; Dixon, P.K.; Liu, A.J.; Durian, D.J. Statistical mechanics of a gas-fluidized particle. *Nature* **2004**, *427*, 521–523. [[CrossRef](#)] [[PubMed](#)]
25. Feitosa, K.; Menon, N. Fluidized granular medium as an instance of the fluctuation theorem. *Phys. Rev. Lett.* **2004**, *92*, 164301. [[CrossRef](#)] [[PubMed](#)]
26. Carberry, D.M.; Reid, J.C.; Wang, G.M.; Sevick, E.M.; Searles, D.J.; Evans, D.J. Fluctuations and irreversibility: An experimental demonstration of a second-law-like theorem using a colloidal particle held in an optical trap. *Phys. Rev. Lett.* **2004**, *92*, 140601. [[CrossRef](#)] [[PubMed](#)]
27. Greinert, N.; Wood, T.; Bartlett, P. Measurement of effective temperatures in an aging colloidal glass. *Phys. Rev. Lett.* **2006**, *97*, 265702. [[CrossRef](#)]
28. Mizuno, D.; Tardin, C.; Schmidt, C.F.; MacKintosh, F.C. Nonequilibrium mechanics of active cytoskeletal networks. *Science* **2007**, *315*, 370–373. [[CrossRef](#)]
29. Gomez-Solano, J.R.; Petrosyan, A.; Ciliberto, S.; Chetrite, R.; Gawędzki, K. Experimental verification of a modified fluctuation-dissipation relation for a micron-sized particle in a nonequilibrium steady state. *Phys. Rev. Lett.* **2009**, *103*, 040601. [[CrossRef](#)]
30. Ciliberto, S. Experiments in stochastic thermodynamics: Short history and perspectives. *Phys. Rev. X* **2017**, *7*, 021051. [[CrossRef](#)]
31. Evans, D.J.; Morriss, G.P. *Statistical Mechanics of Nonequilibrium Liquids*, 1st ed.; Academic Press: London, UK, 1990.
32. Searles, D.J.; Evans, D.J. Ensemble dependence of the transient fluctuation theorem. *J. Chem. Phys.* **2000**, *113*, 3503–3509. [[CrossRef](#)]
33. Evans, D.J.; Searles, D.J. The Fluctuation Theorem. *Adv. Phys.* **2002**, *51*, 1529–1585. [[CrossRef](#)]
34. Evans, D.J.; Searles, D.J.; Williams, S.R. On the fluctuation theorem for the dissipation function and its connection with response theory. *J. Chem. Phys.* **2008**, *128*, 014504. [[CrossRef](#)]

35. Petravić, J.; Evans, D.J. Nonlinear Response for Time-Dependent External Fields. *Phys. Rev. Lett.* **1997**, *78*, 1199–1202. [[CrossRef](#)]
36. Petravić, J.; Evans, D.J. Nonlinear response for nonautonomous systems. *Phys. Rev. E* **1997**, *56*, 1207–1217. [[CrossRef](#)]
37. Todd, B.D. Nonlinear response theory for time-periodic elongational flows. *Phys. Rev. E* **1998**, *58*, 4587–4594. [[CrossRef](#)]
38. Jepps, O.G.; Rondoni, L. A dynamical-systems interpretation of the dissipation function, T-mixing and their relation to thermodynamic relaxation. *J. Phys. A Math. Theor.* **2016**, *49*, 154002. [[CrossRef](#)]
39. Greppi, E.; Rondoni, L. Quantum Exact Response Theory Based on the Dissipation Function. *Entropy* **2025**, *27*, 527. [[CrossRef](#)]
40. Amadori, D.; Colangeli, M.; Correa, A.; Rondoni, L. Exact response theory and Kuramoto dynamics. *Phys. D Nonlinear Phenom.* **2022**, *429*, 133076. [[CrossRef](#)]
41. van Kampen, N. *Stochastic Processes in Physics and Chemistry*, 3rd ed.; North-Holland Personal Library, North-Holland: Amsterdam, The Netherlands, 2007. [[CrossRef](#)]
42. Gardiner, C.W. *Stochastic Methods: A Handbook for the Natural and Social Sciences*, 4th ed.; Springer Series in Synergetics; Springer: Berlin/Heidelberg, Germany, 2009. [[CrossRef](#)]
43. Gallavotti, G. *Nonequilibrium and Irreversibility*, 2nd ed.; Lecture Notes in Physics; Springer: Cham, Switzerland, 2025; Volume 1040. [[CrossRef](#)]
44. Kreuzer, H.J. *Nonequilibrium Thermodynamics and Its Statistical Foundations*, 1st ed.; Monographs on the Physics and Chemistry of Materials; Clarendon Press/Oxford University Press: Oxford, UK; New York, NY, USA, 1981.
45. Spohn, H. *Large Scale Dynamics of Interacting Particles*; Theoretical and Mathematical Physics; Springer: Berlin/Heidelberg, Germany, 1991. [[CrossRef](#)]
46. Colangeli, M.; Rondoni, L. Equilibrium, fluctuation relations and transport for irreversible deterministic dynamics. *Phys. D Nonlinear Phenom.* **2012**, *241*, 681–691. [[CrossRef](#)]
47. Risken, H. *The Fokker-Planck Equation*; Springer: Berlin, Germany, 1996. [[CrossRef](#)]
48. Pavliotis, G.A. *Stochastic Processes and Applications. Diffusion Processes, the Fokker-Planck and Langevin Equations*; Springer: Berlin/Heidelberg, Germany, 2014. [[CrossRef](#)]
49. Balakrishnan, V. *Elements of Nonequilibrium Statistical Mechanics*, 1st ed.; Springer Nature: Cham, Switzerland, 2020; p. 314. [[CrossRef](#)]
50. Colangeli, M.; Duong, M.H.; Muntean, A. A hybrid approach to model reduction of Generalized Langevin Dynamics. *J. Stat. Phys.* **2025**, *192*, 1–20. [[CrossRef](#)]
51. Jones, C.K.R.T. Geometric singular perturbation theory. In *Dynamical Systems: Lectures Given at the 2nd Session of the Centro Internazionale Matematico Estivo (C.I.M.E.) Held in Montecatini Terme, Italy, June 13–22. 1994*; Johnson, R., Ed.; Springer: Berlin/Heidelberg, Germany, 1995; pp. 44–118. [[CrossRef](#)]
52. Sekimoto, K. *Stochastic Energetics*; Lecture Notes in Physics; Springer: Berlin/Heidelberg, Germany, 2010; Volume 799. [[CrossRef](#)]
53. Lepri, S.; Livi, R.; Politi, A. Thermal conduction in classical low-dimensional lattices. *Phys. Rep.* **2003**, *377*, 1–80. [[CrossRef](#)]
54. Chandrasekhar, S. Stochastic Problems in Physics and Astronomy. *Rev. Mod. Phys.* **1943**, *15*, 1. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.