

REVIEW

Open Access



Biomass from microalgae: the potential of domestication towards sustainable biofactories

Manuel Benedetti, Valeria Vecchi, Simone Barera and Luca Dall'Osto* 

Abstract

Interest in bulk biomass from microalgae, for the extraction of high-value nutraceuticals, bio-products, animal feed and as a source of renewable fuels, is high. Advantages of microalgal vs. plant biomass production include higher yield, use of non-arable land, recovery of nutrients from wastewater, efficient carbon capture and faster development of new domesticated strains. Moreover, adaptation to a wide range of environmental conditions evolved a great genetic diversity within this polyphyletic group, making microalgae a rich source of interesting and useful metabolites. Microalgae have the potential to satisfy many global demands; however, realization of this potential requires a decrease of the current production costs. Average productivity of the most common industrial strains is far lower than maximal theoretical estimations, suggesting that identification of factors limiting biomass yield and removing bottlenecks are pivotal in domestication strategies aimed to make algal-derived bio-products profitable on the industrial scale. In particular, the light-to-biomass conversion efficiency represents a major constraint to finally fill the gap between theoretical and industrial productivity. In this respect, recent results suggest that significant yield enhancement is feasible. Full realization of this potential requires further advances in cultivation techniques, together with genetic manipulation of both algal physiology and metabolic networks, to maximize the efficiency with which solar energy is converted into biomass and bio-products. In this review, we draft the molecular events of photosynthesis which regulate the conversion of light into biomass, and discuss how these can be targeted to enhance productivity through mutagenesis, strain selection or genetic engineering. We outline major successes reached, and promising strategies to achieving significant contributions to future microalgae-based biotechnology.

Keywords: Microalgae, Photosynthesis, Photobioreactor, Biomass, Bio-based products, Light-use efficiency, Strain domestication, Molecular genetic

Background

Microalgae, a promising feedstock option

Approximately 100,000 terawatts-year (TW-y) power from sunlight reach the surface of our planet. This is a renewable resource exceeding the current human global energy demand (15 TW-y) and the 24 TW-year predicted for anthropic activities by 2030 [1, 2] by > 3 orders of magnitude. Sunlight might fully provide for future world energy demand [3] and yet its dilute nature represents a major challenge for concentrating, harvesting,

storing it efficiently. Oxygenic photosynthesis converts CO₂ into reduced carbon compounds using light and water; through this process photoautotrophic organisms, namely plants, algae and cyanobacteria, store solar energy at a rate of 120 TW-y at the global scale. Therefore, using sunlight and CO₂ to produce a variety of organic molecules and biomass, by the extensive cultivation of photosynthetic organisms, has the potential to cover a significant portion of global energy demand [4], besides providing for an effective CO₂ capture from e.g. power plants or other large-scale emission sources. As unique feature, photosynthesis allows for direct energy storage into liquid fuels which can be used in the existing

*Correspondence: luca.dallost@univr.it
Dipartimento di Biotecnologie, Università di Verona, Strada Le Grazie 15,
37134 Verona, Italy



transport systems, while other forms of renewable energy such as hydro-, wind or photovoltaic power, cannot.

Thus, mass culture of microalgae has gained interest in the past few decades. Indeed, beside small-scale traditional cultivations mainly aimed to human feeding, commercial production of algae on a larger scale has been identified in recent years as a renewable and environmentally sustainable strategy for feedstock production. Microalgae include a wide group of photosynthetic, eukaryotic, unicellular organisms: green microalgae, belonging to the class Chlorophyceae, include genera which are among the most widely used for industrial applications, such as *Haematococcus*, *Chlorella* and *Dunaliella*. Diatoms and cyanobacteria, which also represent a valuable biotechnological platform [5, 6], will not be included in this review.

The phyletic group of green microalgae include species which have adapted to diverse environmental conditions, even extreme, of the planet [7]. By considering that the unclassified species likely represent the majority of this group [8], it comes that green microalgae are a source of metabolic and genetic diversity [9]. Microalgal biomass represents an energy-rich feedstock, which received increasing attention for commercial cultivation in open ponds or closed photobioreactors (PBRs). So far industrial applications include production of bioactive compounds [10], recombinant proteins [11], next generation biofuels and wastewater treatment [12]. Once target products extracted, the residual biomass can be further processed into livestock feed, organic fertilizer and biostimulants, or used for energy cogeneration [13–15]; therefore, biorefinery processes applied to mono-species cultivation can yield a large variety of resources.

Although photosynthetic machinery is similar to that of plants, microalgae convert solar energy into biomass and fix CO₂ at efficiencies that are appreciably higher than land plants [16]. The maximal conversion efficiencies of solar radiation into biomass are 4.6% for C3 plants and 6.0% for C4 plants at 30 °C, which drops to 2.9% and 4.2% respectively, when measured in the field [17].

Theoretical quantification of 8–10% in energy conversion efficiency of microalgae [18] translates into an expected maximal productivity of 280 ton of algal biomass hectare (ha)⁻¹ year⁻¹, while outdoor mass cultivation record yield beyond 100 ton ha⁻¹ year⁻¹ could not be reached [19]. This compares to 0.2% of energy conversion efficiency and an average of 10 ton ha⁻¹ year⁻¹, reported for sugarcane field trials in the tropics. When considering oil yield extracted from plant vs. algal biomass, palm oil can produce a maximum of 4–5 ton ha⁻¹ year⁻¹ [20, 21] vs. 30 ton ha⁻¹ year⁻¹ [19]. Thus, record yields of microalgal culture at temperate latitudes is >5 times higher respect to the case of the best figure for a plant crop.

Multiple reasons contribute to such a feature:

- i. When growing in aerated liquid cultures, cells have easy access to light, CO₂ and nutrients. They lack non-productive (heterotrophic) organs to maintain, and the simpler unicellular structure make the whole biomass fully photosynthetically active, irrespective to seasonal life cycle;
- ii. Algae are metabolically flexible and have a short doubling time. Although most microalgae are primarily photoautotrophs, many species undergo metabolic shift to heterotrophy upon changes in environmental conditions, utilizing organic compounds as C and energy source or to mixotrophy (carrying out photosynthesis as the main energy source, and both organic molecules and carbon dioxide are used as C source). Moreover, under optimal growth conditions, most species have doubling time of a few hours, and cultures reach as much as 10 g l⁻¹ of heterotrophic dry weight (DW) biomass and 6 g l⁻¹ of photoautotrophic DW biomass [22, 23];
- iii. Microalgae do not require fertile land, and can grow in wastelands, using brackish or waste water, or even sea water in the case of marine species; hence, their cultivation does not compete with resources for conventional food production, and would be more environmentally sustainable respect to extensive cultivation of crops. Therefore, microalgae offer the opportunity to shift part of unsustainable farming and fishing routines toward unproductive region;
- iv. Different species can be selected for specific growth conditions, suited to the local climate, which is more difficult with conventional crops.

Main text

The most promising microalgae species for production of valuable compounds and for biotechnology applications

In this paragraph, the bio-technological applications of best wild type species are reviewed with focus on high-value production chemicals and biomass for biofuels (Table 1). Currently, the most relevant microalgal species for high-value chemicals production are the cyanobacterium *Arthrospira platensis* (formerly known as *Spirulina*) and the green microalgae *Chlorella vulgaris*, *Dunaliella salina* and *Haematococcus pluvialis*, which are mainly dedicated to the production of single products in large-scale cultivation systems.

Arthrospira platensis is exploited as source of nutraceuticals [24], long-chain polyunsaturated fatty acids (lc-PUFAs) [25], carotenoids [26] and proteins [27].

Table 1 Noteworthy microalgae species and their biotechnological applications for production of high-value metabolites

Application	Industrial and medical field	Cosmetics and food colorant	High-value metabolites	Biofuel	Dietary supplement and nutraceuticals
Bioproduct category Microalgae species (metabolite)	Polysaccharides <i>Chlorella</i> spp. (β -glucans, starch) <i>Porphyridium cruentum</i> <i>Netrium digitus</i> Phycotoxins <i>Amphidinium</i> <i>Dinophysis</i> <i>Prorocentrum</i> Phycobilliproteins Red algae (Phycocerythrin)	Phycobilliproteins <i>Arthrospyra platensis</i> (phycocyanin) Carotenoids <i>Arthrospyra platensis</i> <i>Chlorella vulgaris</i> <i>Haematococcus pluvialis</i> <i>Chlorella zofingiensis</i> (astaxanthin) <i>Dunaliella salina</i> (β -carotene)	Mycosporine-like amino acids <i>Aphanizomenon flos-aquae</i> Vitamins <i>Euglena gracilis</i> (biotin, α -tocopherol) <i>Prototheca moriformis</i> (ascorbic acid) <i>Arthrospyra platensis</i> <i>Chlorella</i> spp. Proteins <i>Arthrospyra platensis</i> <i>Chlorella</i> spp.	Oil to biodiesel <i>Botryococcus braunii</i> <i>Chlorella</i> spp. <i>Dunaliella salina</i> <i>Monoraphidium contortum</i> <i>Scenedesmus</i> spp. Carbohydrate to bioethanol <i>Spirogyra</i> spp. <i>Chlorococum</i> spp. Bio-hydrogen <i>Chlamydomonas reinhardtii</i>	Polyunsaturated fatty acids (PUFAs) <i>Parietochloris incise</i> <i>Porphyridium</i> spp. (arachidonic acid) <i>Arthrospyra platensis</i> <i>Rhodomonas salina</i> <i>Tetraselmis uecica</i> (α -linolenic acid) <i>Chlorella minutissima</i> <i>Monodosus</i> spp. <i>Nannochloropsis</i> spp. <i>Neochloris oleoabundans</i> <i>Pavlova lutheri</i> (eicosapentaenoic acid) <i>Cryptocodium</i> spp. <i>Isochrysis galbana</i> <i>Schizochytrium</i> spp. <i>Thalassiosira</i> spp. <i>Thraustochytrium</i> spp. (docosahexaenoic acid)

Other applications of *Spirulina* are in the medical field as a therapeutic [28, 29], as antioxidant [25] and for the extraction of the blue pigment phycocyanin, approved as food colorant by FDA [30].

Genus *Chlorella* includes a number of species which are widely commercialized for production of nutraceuticals. Besides to the high protein, carotenoids and vitamins content [31], *C. vulgaris* contains also β - and α -glucans, D-glucose polysaccharides which act as immune stimulators, free-radical scavengers and anti-cancer compounds [32]. Moreover, *Chlorella* has been successfully used to produce starch, reaching 26% DW yield under mixotrophic condition [33].

Carotenoids represent the commercial product from microalgae with highest success. Carotenoids are widely used as food colorants, aquaculture feed additives and components for cosmetics and skin care; carotenoids also have biomedical applications, including anti-inflammatory activities which are related to their strong antioxidant properties [34]. β -carotene, the first carotenoid successfully marketed at large scale, is produced from the halophilic alga *Dunaliella salina* through both extensive cultivation in ponds and intensive cultivation in PBRs [35]. Recently, new strains with different ability to accumulate carotenoids and different capacity of photoprotection against high light stress have been isolated; the most promising strain was characterized by a β -carotene productivity of $3.5 \text{ g l}^{-1} \text{ day}^{-1}$ at $1500 \mu\text{mol m}^{-2} \text{ s}^{-1}$ [36]. Currently, various strains of *D. salina* growing at different salinity conditions are available [37].

Astaxanthin is a high-value, red keto-carotenoid, successfully commercialized by many companies worldwide through cultivation of the green alga *Haematococcus pluvialis*. Under various stress conditions, this alga changes from a thin-wall mobile phase to a red thick-wall resting phase, in which astaxanthin can reach up to 5% DW [38]. Astaxanthin is widely employed in the feed, cosmetic, aquaculture, nutraceutical and pharmaceutical industries because of its antioxidant potential [39]. Moreover, astaxanthin-rich *Haematococcus* is a popular nutraceutical antioxidant for human diet [40]. *Chlorella zofingiensis* has been proposed as an alternative astaxanthin source which is more reliable in growth [41].

Biosynthesis of fatty acids and triglycerides (TAGs) is relevant for several industrial applications. Microalgae are the primary producers of lc-PUFAs such as eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), which accumulate in the oil of many fish species. Aquaculture farming increased demand of lc-PUFAs for nutrition, which are currently produced from fish oil, while a more sustainable lc-PUFA supply is sought. Several marine algal species are rich in lc-PUFAs thus have a great potential for biorefinery: these include *Thraustochytrium* sp., *Pavlova lutheri*, *Nannochloropsis gaditana*, *Isochrysis galbana*, *Cryptocodium cohnii* (rich in DHA and EPA), *Rhodomonas salina* and *Tetraselmis suecica* (α -linolenic acid) [42, 43] and *Parietochloris incisa* (arachidonic acid) [44]; these are currently exploited by a number of small companies, marketing biomass of high-values but at small-scale. Lc-PUFAs are important elements for human diet:

DHA plays a crucial role as anti-inflammation molecule in allergic diseases and has considerable benefits on visual and cognitive functions; optimization of the ratio of lc-PUFAs in nutraceuticals may contribute to reduce the severity of allergic disease symptoms [45]; oil from *Nannochloropsis*, *Rhodomonas* and *Tetraselmis* has higher antioxidant properties respect to fish oil, likely due to the content of valuable carotenoids and polyphenols and is expected to replace fish oils in diets soon [46].

Further species with high potential for large-scale exploitation of extracellular polysaccharides include the microalgae *Porphyridium* and *Desmidiiales* spp. Red microalgae *Porphyridium* spp. are fast-growing and accumulate extracellular polysaccharides commercially used in cosmetic and medical field [47, 48]. Amongst *Desmidiiales*, *Netrium digitus* has been successfully cultivated in porous substrate bioreactor, reaching a maximum product concentration of 25 g m⁻² [49].

In the last decade, microalgae have received increasing interest as a source of biomass for replacing fossil fuels. Liquid fuels derived from raw biomass are an attractive source of renewable energy, to be used in transport system or energy cogeneration. With respect to the major biofuels currently produced worldwide, namely bio-ethanol from sugar cane and biodiesel from oil crops, the so called “third generation” (microalgal) biofuels, are considered as a promising option, since these organisms are highly productive and provide a solution for the food vs. fuel problem [50].

Green algae accumulate high levels of polysaccharides both as cell-wall constituents and storage molecules that can be fermented to bioethanol [51, 52]. Oil fraction of algal biomass, which range from 20 to 60% DW depending on the species and growth conditions [19, 53], is processed by transesterification to produce biodiesel. In this respect, promising species belong to the genus *Chlorella* [54], *Scenedesmus* [55] and *Monoraphidium*, the latter showing high productivity and high-quality lipid profile [56]. An unusually rich source of TAGs is the green microalga *Botryococcus braunii* (hydrocarbons constituting up to 75% of its DW), however its potential is limited by the slow growth [57].

Despite great advantages offered by microalgae exploitation, production of third generation biodiesel is still far from being commercially viable [58]. As an alternative, biogas generation through anaerobic digestion of microalgal biomass has been proposed as a more energetically-favorable process [59]. The efficiency of biogas production is species-dependent because is based on cell wall degradability and sometime limited by the content in molecules inhibiting growth of methanogenic *Archaea* [60]. A number of pre-treatment procedures have been

tested, including cell wall disruption by chemical/physical methods or enzymatic hydrolysis, which enhanced bio-methane yield [14, 61, 62].

Finally, microalgae and cyanobacteria can produce bio-hydrogen through photo-fermentation, in an anaerobic process involving oxidation of ferredoxin by the hydrogenase enzyme [63, 64]. Although biological H₂ shows great promise for generating future, large scale sustainable energy, a number of bottlenecks still limit its production [65]; however, recent result [66] identified in *C. reinhardtii* promising targets for genetic engineering of H₂ production capacity while the use of temperature-sensitive conditional PSII mutants has been proposed in order to separate the oxygenic biomass-accumulating phase from the oxygen sensitive hydrogenase activity [67].

Microalgae are gaining importance in the biological offset of polluted matrix, because of their ability to thrive under extreme or polluted condition: they serve for direct carbon capture, a way for reducing CO₂ released by large-scale emission plants [68]. Promising species include *Scenedesmus obliquus*, *Chlorella vulgaris*, *Chlorella protothecoides* and *Spirulina* spp., which can grow up to 15–18% CO₂, although highest productivity was observed around 10% CO₂ [69, 70]. In last years, *N. gaditana* is arising as promising species for CO₂ removal due to a high biofixation rate—more than 1.7 g⁻¹ l⁻¹ day⁻¹ [71].

Growth in open ponds is an established technology for bioremediation of wastewater and nutrient recovery in the form of biomass [12]. The effectiveness of microalgae to use inorganic N and P to sustain growth as well as their capacity to sequester heavy metals and toxic compounds, have been demonstrated with a wide range of wastewaters, and at a range of scales [72, 73].

Technical challenges to cost-effective, large-scale microalgae production

Despite a number of industrial applications of microalgae have been proposed and studied in lab-scale, the only successful commercial exploitation of microalgal mass culture is the production of carotenoids, namely β-carotene by *D. salina*, and astaxanthin by *H. pluvialis* [10]. Other species (*Chlorella* spp., *Spirulina* spp.) produce high-value compounds, however these productions are currently applied at small-scale cultivations, which need significant reduction in operating costs to become competitive with the same molecules extracted from other feedstocks [74]. A number of species have been identified as promising targets for biorefinery approaches [75], which, however, have not yet come to economic viability. Thus, while microalgae represent a promising source of valuable bio-based products, (1) optimization of both cultivation and processing technologies, together

with (2) selection of candidates with high growth rate and cell density, are required to make the process profitable [76]. An overview of the approaches and the major challenges related to point (1) are presented as follows.

In algal biomass pipeline, there are many elements which contribute to the overall cost of the process. Major factors to take into account are (i) the choice of production system, (ii) the strategies to supply nutrients, aeration, mixing, and (iii) how to harvest and process biomass, and (iv) the procedure to avoid infections and contaminants. Different approaches are available, each having benefits and limitations [77].

Microalgae are mainly cultivated in open ponds, which are cheaper to build, and easier to operate and to scale-up than closed systems. Generally, ponds are raceways at depth of 20–30 cm, in which biomass is mixed by paddles or left unstirred. Drawbacks of these systems include the complication of controlling contaminations, and the difficult of keeping constant growth parameters (e.g. temperature, pH, light); moreover, they suffer of low productivities ($<20 \text{ g m}^{-2} \text{ day}^{-1}$) due to poor gas exchange and dark zone, therefore low cell density forces to cover extensive areas and requires high costs for harvesting the biomass.

As an alternative to open ponds, closed PBRs allow for higher productivities ($\sim 0.8\text{--}1.5 \text{ g l}^{-1} \text{ day}^{-1}$ [19], up to 10 times higher than ponds). Lab-scale PBRs include flat reactors, tubular reactors or vertical plastic bags, with more control over the growth environment, in which biomass is mixed by air-lift or by pumping. These configurations allow higher cell density than ponds, thus improve economic viability of production, however they (i) have high building costs and are difficult to scale-up, (ii) can operate in a sterile mode, which however adds to the management fees, and (iii) require a high energy input for gas exchange [78, 79]. Together with light, CO_2 and nutrients must be supplied to maximize the growth rate, and it significantly affects the economy balance. CO_2 can be delivered through direct bubbling, and its distribution in the culture represents an additional cost factor; another challenge is the removal of excess O_2 which, above air level, inhibits photosynthesis [80].

A third production method is the surface-attached algal biofilm, which showed greater yield than suspended culture, and lower land and water requirements, in lab-scale trials. Algal biofilm system thus appears a good option for low-cost productions [81, 82], however more research is needed to move from bench-scale to pilot plant.

Following growth, biomass must be (i) harvested and (ii) processed to dryness. Both steps remain a major obstacle to industrial scale processing and contribute to $\sim 1/3$ of the final biomass cost. Current harvesting methods include chemical, mechanical and bio-based

procedures: electrolytes or polymers are added to flocculate cells; centrifugation or flow filtration are rapid methods, which however implies high investments and operating costs; biological-based methods include auto-flocculation (at high pH, in excess of Ca^{2+} ions), bio-flocculation (caused by secreted polymers) or microbial-induced flocculation [83]. Dewatering and drying of biomass is required, and the low biomass concentration (0.1–1% w/w) affects the cost of the final product [84]. In case the release of the products from cells is required, it should be as more energy-efficient as possible, avoiding the use of expensive solvents, and costs for treating biomass should be minimized. Novel approaches which limit the use of solvents, e.g. based on enzymatic hydrolysis of the cell wall [51, 61], still suffer for expensive enzyme production. To significantly reduce energy penalty of the production process, the waste biomass can be directed to anaerobic digestion and production of biogas, fertilizers, soil amendments or feeds [13, 62, 85].


Figure 1 shows the different stages in the production of algal biomass, including the factors to be considered and optimized, which contribute to the price of bio-products. In consideration of the processing costs at the present state of technology, engineering optimization is necessary to find new, cost-effective methods of producing large quantities of feedstock. However, integration of innovative technical solutions with strains improved by biotechnological approaches, appears essential.

The overall cost of a biomolecule is the results of cellular content of the desired product, and growth rate of the culture, the latter being dependent on the efficiency at which photosynthetically active radiation (PAR) is used to drive photosynthesis. Indeed, an area of promising research aims at improving the light-to-biomass conversion efficiency under mass culture conditions.

Biological constraints in light-to-biomass conversion efficiency

Calculations in [18] provided both theoretical maxima of solar energy conversion efficiencies in photosynthesis and productivity yield of microalgae, equal to 8–10% solar-to-biomass and 280 ton of dry biomass $\text{ha}^{-1} \text{ year}^{-1}$, respectively. Instead, outdoor mass cultivation showed that, with the present technology and wild type strains, annual productivities beyond 80–100 ton DW $\text{ha}^{-1} \text{ year}^{-1}$ cannot be maintained at large scale and over long periods [19]. Overcoming this gap, which limits exploitation of microalgae to their full potential, is therefore essential.

Wild type algal strains suffer of light use inefficiency. Enhancing light-to-biomass conversion efficiency will help counterbalancing the cost of energy and nutrients used in the cultivation system, as well as reducing the




costs of downstream biomass processing, making it a target for genetic improvement.

Regulation of light harvesting capacity is crucial for cells in order to balance light reactions and downstream biochemical events of photosynthesis. Indeed, autotrophs have evolved regulatory mechanisms, to fine-tune continuous transitions between “conservative” and “dissipative” state of absorbed energy. In particular, photosynthesis typically displays a light saturation curve (Fig. 2), in which 3 distinct phases can be identified: (1) at low irradiance, namely when light is the limiting factor, the photosynthetic rate increases linearly with light intensity; (2) at increasing irradiances, the limiting factor becomes CO₂ fixation rate, thus photosynthetic rate increases non-linearly as a function of light; (3) when light intensity overcomes the rate of downstream biochemical reactions, photosystems get rid of energy absorbed in excess, and in this phase photosynthetic rate reaches a plateau.

In a dilute culture of *C. vulgaris*, where light attenuation is minimized, light saturation is reached at around 1200 μmol photons m⁻² s⁻¹. At this irradiance, algae protect themselves from excess illumination by triggering the non-photochemical quenching (NPQ) mechanism,

a feedback-regulated de-excitation of Chls that operates in the PSII, to prevent over-excitation of reaction centers (Fig. 2). Although light-dependent energy quenching is a property of all photosynthetic organisms, large differences in amplitude and kinetics can be observed. Some microalgae, such as *C. zofingiensis*, exhibit constitutively high energy quenching activity [86], while in other species (e.g. *C. reinhardtii*) energy quenching is significantly activated only upon acclimation to excess light conditions [87]. NPQ activates when light excitation flux exceeds CO₂ fixation rate. However, algae can experience very high light intensity, saturating photoprotective mechanisms. Light in excess of photosynthesis saturation level is dissipated rather than contributing to biomass accumulation, or even causes synthesis of reactive oxygen species (ROS), which damage the photosynthetic machinery and affect biomass yield. *C. reinhardtii* mutant *npq4*, devoid of NPQ response, was indeed more susceptible to photo-oxidation [87].

Due to the absorption of pigments bound to the large antenna systems in both photosystems, light distribution within the culture is inhomogeneous, and this strongly contributes to the gap between theoretical vs. real



productivity. Such large arrays of antenna complexes have been selected by evolution as favorable trait, since they maximize the light-harvesting capacity and therefore the adaptation to a natural water environment where light is often scarce and limits growth, and cell density remains low. Contrary to the natural environment, growth conditions in mass cultures requires high cell biomass per volume of installed facility. However, this results in high optical density and light shortage in the deeper layers of the culture while cells at the surface layers intercept most photons, resulting in saturation of photosynthesis, dissipation of excess energy and/or photoinhibition. The most inner layers easily drop below the compensation point of photosynthesis while active respiration consumes energy. Thus, dense algal cultures suffer both photo-deprivation and photo-inhibition, decreasing the overall light-to-biomass conversion efficiency far below the theoretical score. Rapid mixing of biomass is often suggested as a solution to light gradients, but it is not: rapid light/dark cycles between dark and over-saturating irradiances have a deleterious effect on biomass yield [88]. Modeling of the light-response curve of photosynthesis in a culture system [89] suggests that optimal setting of OD in the

culture limits shading while maximizes light absorption and net photosynthesis.

One additional factor which contributes to the inefficiency of photosynthesis is carbon fixation: the enzyme RuBisCO, which catalyzes the carboxylation of Ribulose 1,5-bisphosphate (RuBP), has low affinity for CO₂ and can also use O₂ to give oxygenated substrates [90], which ultimately results in ATP/NADPH consumption and loss of fixed C. To compensate for this, RuBisCO accumulates to as much as 50% of total soluble proteins of the cell. Inefficient light use in mass culture, due to inhomogeneous light distribution, results from limitations in the turnover rate of the Calvin-Benson cycle. The selection of strains with higher RuBP carboxylation activity would therefore be a major goal for the optimization of photosynthesis. Possible strategies include the heterologous expression of variants of RuBisCO with a higher specific activity, or the control of allosteric regulators (e.g. RuBisCO activase) to yield into suppression of oxygenase activity.

Interestingly, many microalgae use biophysical carbon concentrating mechanisms for active retention of inorganic C [91] to increase CO₂ availability at the RuBisCO active site within the pyrenoid, a micro-compartment of

the chloroplast. Engineering such a mechanism into algal species devoid of pyrenoids might augment the overall C fixation rate and thus photosynthetic efficiency.

In conclusion, results reported in this section suggest that modulation of photosynthetic reactions is a key factor controlling biomass yield at both saturating and sub-saturating irradiances, that is worth to be considered for a domestication strategy aimed at improving performances in PBRs.

Promises of domestication by forward genetic in improving photosynthetic efficiency

According to the previous section, a gap between theoretical and real biomass productivities of microalgae originates from the high OD of cells. With respect to the problem of inhomogeneous light distribution, high density planting of crops is a condition limiting PAR penetration thus productivity, and it is not different than the condition of elevated cell density reachable in a PBR. Moreover, biomass production with wild type algal strains is poorly viable likely as farming with ancestral crop varieties.

The so called 'Green Revolution' of agriculture, a domestication based on breeding and phenotypic selection, succeeded in pursuing crop productivities over the past 50 years [92, 93]. Industrial application of microalgae may take advantage of a domestication approach, analogous to that carried out for modern crops. Thus, selection of strains carrying desired traits, together with implementing new alleles by random mutagenesis or genetic engineering, might improve performances in PBRs.

Random mutagenesis is recognized as a powerful technology in mutation breeding, widely employed for strain improvement and for studying the molecular basis of metabolic processes. Forward genetic approach is of particular relevance for algal biotechnology, since it avoids restrictions to GMO for outdoor production system [94]. The most common method for generating genetic variability in a population of microalgae is the mutagenesis induced by either physical methods (UV-light, γ - and X-rays) or chemical mutagens, e.g. *N*'-nitro-*N* nitroso-guanidine (NTG) and ethyl methanesulfonate (EMS).

Attempts for algae genetic improvement, aimed to enhance light-to-biomass conversion efficiency, relied on random mutagenesis and screening of favorable traits. These approaches, while overcame scarcity of genetic engineering tools in microalgae, needed for efficient screening strategies for strains with higher productivity. Some of these approaches, which succeeded in increasing photosynthetic yield, are presented as follows.

Due to detrimental effect of high OD for mass cultivation, strains carrying truncated antenna size were

proposed to perform better in light transmittance than wild type [95]. Mutagenesis and screening of *C. reinhardtii* was employed to isolate mutants having a truncated light-harvesting system [96–100]: all of them showed a higher productivity than the wild type in bench-scale growth systems. Cazzaniga and collaborators [101] applied random mutagenesis to a thermotolerant, fast-growing strain of *C. sorokiniana*, and selected pale-green mutants by imaging Chl fluorescence. Mutants were able to perform photosynthesis more efficiently than wild type, minimizing photoinhibition in high light; the positive effect on photosynthetic productivity was confirmed in both lab-scale and outdoor PBRs. Similar results were obtained with *N. gaditana* strains having reduced cellular pigment content [102]. Finally, simultaneous knock-down of three light-harvesting complex proteins (LHCMB1, 2 and 3) in *C. reinhardtii*, by an RNAi triple knock-down strategy, resulted in improved light-to- H_2 (+180% than wild type) and light-to-biomass (+165%) conversion efficiencies [103].

Implementation of biosynthetic pathway of accessory pigments, e.g. phycobilins or Chls, into genus of industrial interest, has been proposed for improving harvesting efficiency over the full PAR spectrum [4]. Recently, the enzyme responsible for the synthesis of Chl *f*, an oxidized form of Chl *a*, has been isolated from the cyanobacterium *C. fritschii* [104]; heterologous expression of Chl *f* synthase succeeded in accumulating this chromophore in *Synechococcus* sp. Since Chl *f* expands the spectral range for photosynthesis by absorbing far red light, its expression in microalgae may confer advantages for mass culture in PBRs, which suffers for detrimental sieve-effects at high cell densities. However, the feasibility of this approach, which assumes a correct binding of the new chromophores into the existing pigment-binding complexes, still await experimental confirmation.

CO_2 fixation rate is a major limiting step in biomass yield, which arise from RuBisCO inefficiency. Genetic engineering of RuBisCO to increase its catalytic activity or to enhance its specificity towards CO_2 have been proposed as ways to overcome these limitations [4]. It is worth noting that the natural diversity of RuBisCO is limited, likely because the interactions which support catalytic activity make most of the isoforms of this enzyme intolerant to mutations [90], indeed attempts to overcome its limitations by directed evolution, had scant success. Although variants of RuBisCO with higher activity have been identified [105, 106] their heterologous expression in algal strains of industrial potential is still missing. Recently, *E. coli*-based screen of new RuBisCO variants obtained by direct evolution allowed the identification of an unexplored subunit interface with potential of increasing CO_2 fixation rate [107]. Site-directed

mutagenesis in such subunit may generate novel variants whose enzymatic characteristics can be subsequently tested in microalgae in terms of enhanced CO₂ fixation. An additional approach may reside in the generation of hybrid RuBisCO by using novel activase isoforms from chemolithoautotroph microorganisms such as *Acidithiobacillus ferrooxidans* [108].

A high photosynthetic efficiency is attainable only in low irradiance and controlled light environments, which allow most absorbed photons can be utilized by the culture; instead in the outdoor, efficiency drops due to fluctuating irradiances, exceeding the photosynthetic capacity. Autotrophs evolved mechanisms for regulating the efficiency of light capture, which can become target of domestication strategies. Several microalgal species e.g. *C. reinhardtii* and *H. pluvialis*, trigger fast phototactic response, for fine-tuning exposure to light. Indeed, phototaxis confers fitness advantage and it is regulated by cytoplasmic redox balance, which in turn is affected by photosynthetic electron transport rate [109]. In the attempt of isolating strains with improved light-use-efficiency, Kim and collaborators [110] analyzed a *C. reinhardtii* mutant population for rapid phototaxis response and identified mutants with enhanced photoautotrophic growth and lipid production, respectively 1.9- and 8.1-fold increases than wild type.

Photosynthetic organisms dynamically regulate the amplitude of NPQ (see “[Biological constraints in light-to-biomass conversion efficiency](#)” section): by balancing amplitudes of light harvesting vs. energy dissipation, they maintain optimal fitness in changing light environment. The slow relaxation rate of NPQ upon high- to low-light transition was considered to reduce the overall conversion efficiency of solar to biomass in microalgae, consistent with recent evidences in plants [111, 112]. Indeed, deletion of the OCP protein, responsible for NPQ response in cyanobacteria, resulted in a 30% higher biomass yield in mass cultures of *Synechocystis* than wild type cells [113]. Random insertional mutagenesis and Targeting Induced Local Lesions IN Genomes (TILLING) approach on *C. reinhardtii*, following by Chl fluorescence imaging screening, has produced mutants specifically devoid of *lhcsr* genes [87, 114]. In their report, [115] proposed that biomass productivity depends on LHCSR protein accumulation: *C. reinhardtii* strains lacking the two *lhcsr3* genes were more productive than wild type, thus confirming down-regulation of NPQ is a strategy for improving light use efficiency in microalgae. Instead, more recently, [116] observed no significant differences in biomass yield between *C. reinhardtii* wild type and the *npq4 lhcsr1* mutant, devoid of all *Lhcsr* isoforms.

Microalgae growing in mass culture experience rapid changes in the irradiance due to cell mixing into the PBR. The amount of time spent in sub-saturating vs. excess light influences the biomass productivity, which is lower in fluctuating light conditions [115] possibly due to the metabolic energy needed to repair photodamage. Hence, improving photosynthetic efficiency in excess light conditions is potentially a major goal for establishing efficient outdoor cultivation. Research efforts aimed to obtain non-GMO algal strain tolerant to excess light, mainly focused on the model alga *C. reinhardtii*. Förster et al. [117] isolated *very high light resistant* (VHL-R) mutations, which allowed near maximal growth rate at irradiances lethal to the control genotype; characterization of these strains reveals they affected the regulatory pathways which modulate photoprotective response, including PSII repair and ROS detoxification. In [118], wild type strain was UV-mutagenized and plated onto medium containing a lethal concentration of the photosensitizer Red Bengal; by this approach, SOR1 was identified as a factor enhancing tolerance to photooxidative stress conditions. Schierenbeck and coworkers [119] performed UV-mutagenesis followed by selection under high irradiance (2000 $\mu\text{mol m}^{-2} \text{s}^{-1}$); the two mutations selected, which both mapped in the putative Light Responsive Signal 1 (LSR1) gene, conferred an improved resistance of cells against exogenous ROS. Recent results concern the isolation of pale-green, singlet oxygen resistant mutant by EMS-mutagenized *C. vulgaris*, which showed biomass yield enhancement by 68% than wild type strain (Dall'Osto et al. unpublished results).

Improving algal biomass productivity by genetic engineering: methods, state of the art and perspectives

Genetic manipulation approaches have the potential to revolutionize industry based on microalgae cultivation. These include transfer of genes isolated from other species to generate strains with desirable commercial traits such as tolerance to excess light and heat stress, resistance to herbivore/pathogen, capacity to outcompete opportunistic organisms, or to express biosynthetic pathways into more productive strains (Fig. 3). Recent progress in genome sequencing, metagenome/metatranscriptome approaches, and genetic manipulation, yielded into significant advancement in microalgal research. In this paragraph, different strategies of genetic engineering, which revealed effective in improving algal productivity, are discussed. Moreover, additional solutions are proposed.

In the last decades, several efforts have been attempted to optimize the transformation efficiency of different microalgae species. Stable transformation was first developed in *C. reinhardtii*: being able to growth both in




Fig. 3 Schematic depiction of the major desirable traits to be either implemented or improved, toward higher productivity of microalgae in mass culture

autotrophic and heterotrophic conditions, as haploid or diploid cells, *Chlamydomonas* was adopted as powerful genetic system for studying different physiological mechanisms. Nuclear transformation of *C. reinhardtii* may be achieved by several methods such as electroporation [120], *Agrobacterium*-mediated transformation [121], silicon carbide whiskers and positively-charged amino-clay nanoparticles [122, 123] and glass beads agitation method [124]. Both electroporation and *Agrobacterium*-mediated transformation also succeeded in transforming algae of economic interest such as *C. vulgaris*, *Neochloris oleoabundans* and *H. pluvialis* [125–127]. The introduction of foreign genes into the nuclear genome of microalgae is generally guided by a random integration event [128]. Nuclear gene expression is frequently subjected to strong silencing mechanisms due both to position effect and to epigenetic phenomena, similar to those of land plants [129]; indeed in microalgae, silenced multicopy transgenes exhibit high levels of DNA methylation as in land plants [130, 131].

In last years, many efforts have been attempted to increase the heterologous expression potential of microalgae. Increased expression of transgenes was obtained

by fusing the sequences encoding the gene of interest and the selection marker in a unique bicistronic RNA [132, 133]. Recourse to a frequent subset of preferred codons results in elevated transcriptional levels, while the use of codons introducing unintended splicing signals negatively affects the overall expression of transgenes [134], thus a codon usage optimization is mandatory to maximize protein yield. To improve the selection of high-level expressing transformants, last-generation expression vectors exploit the 2A peptide [135] to generate transcriptional fusions between selection marker sequence (e.g. antibiotic resistance) and the gene of interest [132]. Transcription factors are gaining increasing attention as key regulators of metabolic pathways, in order to enhance yield of high-value molecules or to maximize the production of foreign proteins in microalgae. Over-expression of NsbZIP1, a transcription factor carrying the basic leucine zipper, resulted in both enhanced growth rate and higher lipid contents in *N. salina* [136].

The biolistic method (particle-gun bombardment) is the elective procedure for chloroplast transformation in microalgae [137]. The introduction of foreign genes into the plastome is guided by site-specific integration

event (i.e. by homologous recombination). Expressing foreign genes in the chloroplast enables to circumvent gene silencing events, which affect the nuclear expression; moreover, it allows for the introduction of operons, encoding several enzymes of a pathway. However, the resulting transformants must undergo several rounds of selection in order to acquire the homoplasmic condition [138].

Further details concerning nuclear and chloroplast expression in *C. reinhardtii* are summarized in Table 2; moreover, many of these aspects have been covered extensively in a recent review [137].

Genome editing technology enables for both gene deletion and gene integration, therefore implementation of these novel genetic tools in microalgae would allow for manipulation of metabolic networks. Recently, a novel approach based on CRISPR–CAS9 genome editing technology have been successfully developed in both the marine diatom *P. tricornutum* [139] and in *C. reinhardtii*, allowing for deletion of specific gene functions [140]; the latter was achieved by a DNA-free CRISPR–Cas9 method and the outcome was the sequential *FTSY* and *ZEP* double-gene knockout, that resulted in improved photosynthetic productivity. Analogous approach was used to abolish the functions encoded by *MAA7*, *CpSRP43* and *ChlM* genes, which led to pale-green mutants [141]. In *C. reinhardtii*, the replacement of Cas9 with the Cpf1 ribonucleoprotein achieved a more efficient homology-directed DNA replacement [142]. However, a common limitation of free-DNA CRISPR–Cas9/Cpf1 methods resides in the lack of selection markers that, in turn, hinders a straightforward selection of the desired mutants; being the genome editing event induced at very low frequency (0.5–0.6%), a visible phenotype makes mutant selection easier. A DNA-based CRISPR–CAS9 method has been developed in the industrial oleaginous microalga *N. oceanica* [143] in which nuclear transformation

can be efficiently performed by introducing expression cassettes obtained by PCR, making unnecessary the use of expression vectors [144]. Recently, a doubling of the lipid production in *N. gaditana* was obtained by deleting a transcription factor that acts as negative regulator in lipid biosynthesis [145].

As previously described, a truncated antenna size yielded into increased productivity in green microalgae (see “Promises of domestication by forward genetic in improving photosynthetic efficiency” section), thus proteins involved in the biogenesis of photosynthetic machinery can be targeted for increasing biomass production. Truncated light-harvesting antenna 1 (*TLA1*), a nuclear gene putatively involved in the regulation of the antenna size of *C. reinhardtii*, was up- and down-regulated by overexpression and RNAi, respectively. The strain over-expressing *TLA1* showed a larger antenna size for both photosystems and lower Chl *a/b* ratio than the wild type, while its down-regulation resulted in the opposite phenotype changes [146]. LHCII, the nucleus-encoded light-harvesting proteins associated with PSII, tunes the light harvesting capacity to the prevailing light condition. In *C. reinhardtii*, LHCII translation efficiency is regulated by the cytosolic RNA-binding protein NAB1, which is subjected to specific nitrosylation in limiting light, thus making such repressor less-active and promoting accumulation of LHC [147].

Manipulation of RuBisCO activity, namely the major constraint for C assimilation e.g. under excess light conditions [148], may improve the photosynthetic yield [149]. Although site-directed mutants in the *rbcl* (RuBisCO large subunit) gene [150, 151] as well as hybrid variants with altered specificity of RuBisCO–RuBisCO activase interaction [152] have been generated, their over-expression in *C. reinhardtii* did not increase the overall photosynthetic yield.

Table 2 Transformation of *C. reinhardtii*

Organelle	DNA-delivery method	Genetic mechanism	Advantages	Disadvantages	Selection marker
Nucleus	Electroporation gene-gun bombardment <i>A. tumefaciens</i> -mediated Glass beads Silicon carbide whiskers and aminoclay nanoparticles	Ectopic recombination (random integration)	Protein can be expressed as secreted protein Post-translational modifications	Identification of high-expressing Transformants gene silencing	Resistance to Paromycin, Zeocin, Hygromycin, Chloramphenicol Auxotrophic complementation (<i>ARG7</i> , <i>NIT1</i> , <i>oeel</i>)
Chloroplast	Gene-gun bombardment Glass beads	Homologous recombination (site-specific integration)	Compartmentalization Lacks gene silencing High expression level	Need to identify homoplasmic transformant Lacks post-translational modifications	resistance to Spectinomycin Auxotrophic complementation (<i>atpB</i> , <i>psbH</i>)

This table displays DNA-delivery methods, genetic mechanism driving the transformation, and the most common selection markers employed so far. Major advantages and disadvantages of nuclear vs. chloroplast transformation are reported

Recently, biomass productivity as well as lipid yield increased up to 40% in the oleaginous *Nannochloropsis oceanica* by overexpressing endogenous RuBisCO activase [153]. Conversely, a reduction in the RuBisCO activity by site directed mutagenesis resulted in a ten-fold higher H₂ production in *C. reinhardtii* [154], likely because Calvin-Benson cycle competes with Hydrogenase for reducing equivalents.

Further strategies included (i) the PCR-based gene shuffling of *Chlamydomonas* rbcL with sequences representing natural variants of this gene, which yielded isoforms with higher V_{\max} of carboxylation catalysis [155]; (ii) regulation of RuBisCO accumulation according to culture conditions, by tuning mRNA level of the nuclear maturation factor MRL1 [156]; (iii) overexpression of sedoheptulose 1,7-bisphosphatase from *C. reinhardtii*, which succeeded in enhancing photosynthetic efficiency in *D. bardawil* [157]; finally, the over-expression of Low-CO₂ Inducible (LCI) proteins in *C. reinhardtii*, under conditions which typically repress their synthesis (i.e. high CO₂ concentration), increased biomass production under elevated CO₂ conditions as much as 80% than control strain [158].

High productivity in open ponds is restricted to species which adapted to high salt concentration (e.g. *Dunaliella*) or high pH (e.g. *Spirulina*), thus outcompeting naturally occurring contaminants. Hence, a trait which confers competitive advantage over undesirable microorganisms, is crucial both to increase the biomass productivity and to reduce the operating costs for maintenance of axenic cultures (Fig. 1), particularly in either open pond or heterotrophic conditions. In this perspective, non-canonical substrates may be employed for sustaining algal growth. The expression of the phosphite dehydrogenase D (PTXD) from *P. stutzeri* WM88 [159] confers to *C. reinhardtii* the capacity of metabolizing phosphite, namely a P source which cannot be utilized by plants, fungi and most bacteria. Transgenic *Chlamydomonas* cells showed higher fitness than *S. obliquus* in competition experiments in which phosphite-repleted/phosphate-depleted medium was used [160].

Some algal species are strict autotrophs or are highly selective for their C source (e.g. *Chlamydomonas* for acetate), thus trophic conversion by metabolic engineering would be desirable. *Chlamydomonas* cells expressing the hexose transporter HUP1 (monosaccharide-H⁺ symporter from *C. kessleri*) metabolized externally supplied glucose for heterotrophic growth and showed higher H₂ production capacity; however, results suggest that glucose cannot fully replace acetate as a C source, for long-term growth in the dark [161].

Other algal species can metabolize a large array of sugars, and strongly increase their productivity under heterotrophic or mixotrophic growth conditions [162]; however, heterotrophic growth requires additional costs due to need for exogenous carbon source and maintenance of axenic conditions. Algal strains able of metabolizing raw lignocellulosic biomass scraps, namely cheap agricultural wastes, would certainly contribute to make the whole process economically viable. Foreign genes encoding bacterial and fungal plant Cell Wall Degrading Enzymes (CWDEs), were constitutively expressed in microalgae and addressed to the secretory pathway [132]. In *C. reinhardtii*, yield of secreted proteins was improved up to eight-fold by fusing both the putative signal peptide of gametolysin and the repeated serine-proline module, to the N and C terminus of the recombinant protein, respectively [163]. Contrary to plant cell, some species of unicellular green algae possess a cell wall mainly constituted by proteins (e.g. *C. reinhardtii*) [164], thus lack of polysaccharides as major components circumvents the deleterious effects of expressing CWDEs in plants, likely related to hyper-immune responses [165, 166]. Although some algal spp. synthesizes endogenous CWDEs [167], the native cellulolytic machinery is not efficient enough for degrading hydrolysis-recalcitrant substrates such as lignocellulose. Thus, a promising perspective is the expression of a range of secreted CWDEs, including polygalacturonases, hemicellulases, cellulases and ligninases in a unique algal culture, analogously to the approaches developed in yeasts which yielded into strains able to grow on cellulosic substrates [168].

An overview of the major genetic manipulations which may lead to an improvement of biomass productivity is represented in Fig. 4.

Conclusions

Commercially cultivated for several decades, microalgae are now recognized to offer a great potential for exploitation in different fields including pharmaceuticals, aquaculture and renewable energies. Former efforts in their industrial applications mainly focused in optimizing culture parameters and selecting the best performing wild type strain. However, to promote cultivation of microalgae as a new biotechnological sector, a number of challenges still have to be overcome. Domestication strategies achieved by genetic and metabolic engineering will be crucially important to isolate “*smart strains*” with improved yield, in order to make the production successfully marketed. The opportunities offered by investments in both basic and applied research, are considerable: (i) the rapid evolution of genome sequencing techniques will help




Fig. 4 Potential traits to be implemented in GM—*C. reinhardtii* cell. The diagram displays a number of genetic strategies, aimed to enhance productivity in mass culture of microalgae. Gene over-expression (OE) using hybrid promoters or viral cis-acting elements and gene disruption/down-regulation (KO/KD) by Crispr–Cas9 and RNAi approaches are indicated. Some traits that may result in higher productivity include an increased photosynthetic efficiency, improved phototaxis, the use of non-canonical substrates, and optimized carotenoid, lipid and isoprene metabolism. Up- and down-ward pointing arrow mean up- and down-regulation, respectively, and are referred to the expression level of the corresponding endogenous enzyme. Bulb and red cross mean enzymatic in vitro improvement and loss of function, respectively. Abbreviations: *Chl-fS* chlorophyll *f* synthase, *CWDE* cell-wall degrading enzyme, *FTSY* chloroplast signal recognition particle, *GL* gametolysin signal peptide, *HS* hydrocarbon-synthase, *HUP1* hexose-proton symporter, *LHC* light harvesting complexes, *ME* malate dehydrogenase, *ML* multifunctional lipase, *NAB1* RNA-binding protein, *PHY D* phytoene desaturase, *PHY S* phytoene synthase, *PS* patchoulol synthase, *PTXD* phosphite dehydrogenase, β -*PS* β -phellandrene synthase, *TF* transcription factor, *TLA1* truncated light-harvesting antenna 1, *ZEP* zeaxanthin epoxidase

defining the gene networks controlling growth, while -omics approaches allow to identify regulatory points of cellular pathways, thus enabling manipulation of key metabolic steps; (ii) prospective redesigns of algal system include light-to-biomass conversion efficiency, oil content/composition, nutrient recovery capacity; (iii) extend the genetic transformation techniques, now carried out successfully in few species only, to the most industrially-relevant species, will offer the opportunity to address the biological constraints limiting growth yield; finally, (iv) the development of reproducible genome editing techniques will permit a fine matching

of the primary metabolism to the mass culture conditions, or the development of molecular strategies for strain containment. Encouraging results have recently been obtained by boosting light-use-efficiency or by strengthening specific metabolic pathways. Additional research efforts and funding for implementing innovative biorefineries, will realistically support progress toward next-generation algal biotechnology.

Abbreviations

Chl: chlorophylls; CWDE: cell wall degrading enzyme; DHA: docosahexaenoic acid; DW: dry weight; EMS: ethyl methanesulfonate; EPA: eicosapentaenoic acid; GMO: genetically-modified organism; lc-PUFA: long-chain

polyunsaturated fatty acid; LHCBM/LHCSR: light-harvesting complex proteins; NGT: *N*-nitro-*N* nitrosoguanidine; NPQ: non-photochemical quenching; OD: optical density; PAR: photosynthetically active radiation; PBR: photobioreactor; PSII: photosystem II; ROS: reactive oxygen species; RuBisCO: ribulose 1,5-bisphosphate carboxylase-oxygenase; RuBP: ribulose 1,5-bisphosphate; TAGs: triglycerides; TW-y: terawatts-year.

Authors' contributions

MB and LD identified patterns and trends in the literature and designed the structure of the review. VV and SB prepared the tables and figures. All of the authors contributed to searching for relevant literature, and carried out a critical analysis of the literature, discussed together and wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We thank Prof. Roberto Bassi for his suggestions made during the preparation of this review. Because of space constraints, a number of noteworthy publications has not been cited or discussed properly. We apologize to the authors for this lack of completeness.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

Not applicable.

Consent for publication

Not applicable.

Ethics approval and consent to participate

Not applicable.

Funding

Work in our lab is supported by grants from the University of Verona (Fondo Ricerca di Base HuntingLight—Grant Number 201190).

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 8 July 2018 Accepted: 31 October 2018

Published online: 10 November 2018

References

- Cho A. Energy's tricky tradeoffs. *Science*. 2010;329:786–7.
- Hambourger M, Moore GF, Kramer DM, Gust D, Moore AL, Moore TA. Biology and technology for photochemical fuel production. *Chem Soc Rev*. 2009;38:25–35.
- Chu S, Majumdar A. Opportunities and challenges for a sustainable energy future. *Nature*. 2012;488:294–303.
- Stephenson PG, Moore CM, Terry MJ, Zubkov MV, Bibby TS. Improving photosynthesis for algal biofuels: toward a green revolution. *Trends Biotechnol*. 2011;29:615–23.
- Hagemann M, Hess WR. Systems and synthetic biology for the biotechnological application of cyanobacteria. *Curr Opin Biotechnol*. 2018;49:94–9.
- Huang W, Daboussi F. Genetic and metabolic engineering in diatoms. *Philos Trans R Soc B Biol Sci*. 2017;372:1728.
- Tirichine L, Bowler C. Decoding algal genomes: tracing back the history of photosynthetic life on earth. *Plant J*. 2011;66:45–57.
- Guiry MD. How many species of algae are there? *J Phycol*. 2012;48:1057–63.
- Blunt JW, Copp BR, Keyzers RA, Munro MHG, Prinsep MR. Marine natural products. *Nat Prod Rep*. 2012;29:144–222.
- Borowitzka MA. High-value products from microalgae—their development and commercialisation. *J Appl Phycol*. 2013;25:743–56.
- Scranton MA, Ostrand JT, Fields FJ, Mayfield SP. Chlamydomonas as a model for biofuels and bio-products production. *Plant J*. 2015;82:523–31.
- Sutherland DL, Howard-Williams C, Turnbull MH, Broady PA, Craggs RJ. Enhancing microalgal photosynthesis and productivity in wastewater treatment high rate algal ponds for biofuel production. *Bioresour Technol*. 2015;184:222–9.
- Dineshkumar R, Kumaravel R, Gopalsamy J, Sikder MNA, Sampathkumar P. Microalgae as bio-fertilizers for rice growth and seed yield productivity. *Waste Biomass Valoriz*. 2018;9:793–800.
- Passos F, Uggetti E, Carrère H, Ferrer I. Pretreatment of microalgae to improve biogas production: a review. *Bioresour Technol*. 2014;172:403–12.
- García-Gonzalez J, Sommerfeld M. Biofertilizer and biostimulant properties of the microalga *Acutodesmus dimorphus*. *J Appl Phycol*. 2016;28:1051–61.
- Bhola V, Swalaha F, Ranjith Kumar R, Singh M, Bux F. Overview of the potential of microalgae for CO₂ sequestration. *Int J Environ Sci Technol*. 2014;11:2103–18.
- Zhu X-G, Long SP, Ort DR. Improving photosynthetic efficiency for greater yield. *Annu Rev Plant Biol*. 2010;61:235–61.
- Melis A. Solar energy conversion efficiencies in photosynthesis: minimizing the chlorophyll antennae to maximize efficiency. *Plant Sci*. 2009;177:272–80.
- Rodolfi L, Zittelli GC, Bassi N, Padovani G, Biondi N, Bonini G, et al. Microalgae for oil: strain selection, induction of lipid synthesis and outdoor mass cultivation in a low-cost photobioreactor. *Biotechnol Bioeng*. 2009;102:100–12.
- Chisti Y. Biodiesel from microalgae. *Biotechnol Adv*. 2007;25:294–306.
- Mata TM, Martins AA, Caetano NS. Microalgae for biodiesel production and other applications: a review. *Renew Sustain Energy Rev*. 2010;14:217–32.
- Mitra D, van Leeuwen (Hans) J, Lamsal B. Heterotrophic/mixotrophic cultivation of oleaginous *Chlorella vulgaris* on industrial co-products. *Algal Res*. 2012;1:40–8.
- Cuaresma M, Janssen M, Vilchez C, Wijffels RH. Productivity of *Chlorella sorokiniana* in a short light-path (SLP) panel photobioreactor under high irradiance. *Biotechnol Bioeng*. 2009;104:352–9.
- Herrera Bravo de Laguna I, Toledo Marante FJ, Luna-Freire KR, Mioso R. Extraction of nutraceuticals from *Spirulina* (blue-green alga): a bioorganic chemistry practice using thin-layer chromatography. *Biochem Mol Biol Educ*. 2015;43:366–9.
- Al-Dhabi NA, Valan Arasu M. Quantification of phytochemicals from commercial spirulina products and their antioxidant activities. *Evid Based Complement Alternat Med*; 2016. Article ID: 7631864.
- Hynstova V, Sterbova D, Klejdus B, Hedbavny J, Huska D, Adam V. Separation, identification and quantification of carotenoids and chlorophylls in dietary supplements containing *Chlorella vulgaris* and *Spirulina platensis* using high performance thin layer chromatography. *J Pharm Biomed Anal*. 2018;148:108–18.
- Lupatini AL, Colla LM, Canan C, Colla E. Potential application of microalga *Spirulina platensis* as a protein source. *J Sci Food Agric*. 2017;97:724–32.
- Wu Q, Liu L, Miron A, Klímová B, Wan D, Kuča K. The antioxidant, immunomodulatory, and anti-inflammatory activities of *Spirulina*: an overview. *Arch Toxicol*. 2016;90:1817–40.
- Lima FAV, Joventino IP, Joventino FP, de Almeida AC, Neves KRT, do Carmo MR, et al. Neuroprotective activities of *Spirulina platensis* in the 6-OHDA model of Parkinson's disease are related to its anti-inflammatory effects. *Neurochem Res*. 2017;42:3390–400.
- Rizzo RF, Santos BNC, Castro GFPS, Passos TS, Nascimento MA, Guerra HD et al. Production of phycobiliproteins by *Arthrospira platensis* under different light conditions for application in food products. *Food Sci Technol*. 2015;35(2):247–52.
- García JL, de Vicente M, Galán B. Microalgae, old sustainable food and fashion nutraceuticals. *Microb Biotechnol*. 2017;10:1017–24.
- Tabarsa M, Shin IS, Lee JH, Surayot U, Park WJ, You SG. An immune-enhancing water-soluble α -glucan from *Chlorella vulgaris* and structural characteristics. *Food Sci Biotechnol*. 2015;24:1933–41.

33. Li T, Gargouri M, Feng J, Park J-J, Gao D, Miao C, et al. Regulation of starch and lipid accumulation in a microalga *Chlorella sorokiniana*. *Bioresour Technol*. 2015;180:250–7.
34. Young AJ, Lowe GL. Carotenoids—antioxidant properties. *Antioxidants*. 2018;7(2):28. <https://doi.org/10.3390/antiox7020028>.
35. Bonnefond H, Moelants N, Talec A, Mayzaud P, Bernard O, Sciandra A. Coupling and uncoupling of triglyceride and beta-carotene production by *Dunaliella salina* under nitrogen limitation and starvation. *Biotechnol Biofuels*. 2017;10:25.
36. Xu Y, Ibrahim I, Wosu C, Ben-Amotz A, Harvey P. Potential of new isolates of *Dunaliella salina* for natural β -carotene production. *Biology*. 2018;7:14.
37. Ahmed RA, He M, Aftab RA, Zheng S, Nagi M, Bakri R, et al. Bioenergy application of *Dunaliella salina* SA 134 grown at various salinity levels for lipid production. *Sci Rep*. 2017;7:8118.
38. Wayama M, Ota S, Matsui H, Nango N, Hirata A, Kawano S. Three-dimensional ultrastructural study of oil and astaxanthin accumulation during encystment in the green alga *Haematococcus pluvialis*. *PLoS ONE*. 2013;8:e53618.
39. Shah MMR, Liang Y, Cheng JJ, Daroch M. Astaxanthin-producing green microalga *Haematococcus pluvialis*: from single cell to high value commercial products. *Front Plant Sci*. 2016;7:531.
40. Nicoletti M. Microalgae nutraceuticals. *Foods*. 2016;5:54.
41. Liu J, Sun Z, Gerken H, Huang J, Jiang Y, Chen F. Genetic engineering of the green alga *Chlorella zofingiensis*: a modified norflurazon-resistant phytoene desaturase gene as a dominant selectable marker. *Appl Microbiol Biotechnol*. 2014;98:5069–79.
42. Liu J, Sommerfeld M, Hu Q. Screening and characterization of *Isochrysis* strains and optimization of culture conditions for docosahexaenoic acid production. *Appl Microbiol Biotechnol*. 2013;97:4785–98.
43. Ji X-J, Ren L-J, Huang H. Omega-3 biotechnology: a green and sustainable process for omega-3 fatty acids production. *Front Bioeng Biotechnol*. 2015;3:158.
44. Zorin B, Grundman O, Khozin-Goldberg I, Leu S, Shapira M, Kaye Y, et al. Development of a nuclear transformation system for oleaginous green alga *Lobosphaera (Parietochloris) incisus* and genetic complementation of a mutant strain, deficient in arachidonic acid biosynthesis. *PLoS ONE*. 2014;9:e105223.
45. Zárate R, el Jaber-Vazdekis N, Tejera N, Pérez JA, Rodríguez C. Significance of long chain polyunsaturated fatty acids in human health. *Clin Transl Med*. 2017;6:25.
46. Ryckebosch E, Bruneel C, Termote-Verhalle R, Goiris K, Muylaert K, Foubert I. Nutritional evaluation of microalgae oils rich in omega-3 long chain polyunsaturated fatty acids as an alternative for fish oil. *Food Chem*. 2014;160:393–400.
47. Arad SM, van Moppes D. Novel sulfated polysaccharides of red microalgae: basics and applications BT-absorption and adsorption of heavy metals by microalgae. *Absorpt adsorpt heavy met by microalgae*. Hoboken: Wiley; 2013. p. 406–16.
48. Dvir I, Stark AH, Arad (Malis) S. A foodomics approach reveals hypocholesterolemic activity of red microalgae. *Genomics, proteomics metabolomics nutraceuticals funct foods*. Hoboken: Wiley; 2015. p. 31–9.
49. Ekelhof A, Melkonian M. Enhanced extracellular polysaccharide production and growth by microalga *Netrium digitus* in a porous substrate bioreactor. *Algal Res*. 2017;28:184–91.
50. Milano J, Ong HC, Masjuki HH, Chong WT, Lam MK, Loh PK, et al. Microalgae biofuels as an alternative to fossil fuel for power generation. *Renew Sustain Energy Rev Pergamon*. 2016;58:180–97.
51. Shokrkar H, Ebrahimi S, Zamani M. Bioethanol production from acidic and enzymatic hydrolysates of mixed microalgae culture. *Fuel*. 2017;200:380–6.
52. Shokrkar H, Ebrahimi S, Zamani M. Enzymatic hydrolysis of microalgal cellulose for bioethanol production, modeling and sensitivity analysis. *Fuel*. 2018;228:30–8.
53. Ghasemi Naghdi F, González González LM, Chan W, Schenk PM. Progress on lipid extraction from wet algal biomass for biodiesel production. *Microb Biotechnol*. 2016;9:718–26.
54. Liu J, Mao X, Zhou W, Guarnieri MT. Simultaneous production of triacylglycerol and high-value carotenoids by the astaxanthin-producing oleaginous green microalga *Chlorella zofingiensis*. *Bioresour Technol*. 2016;214:319–27.
55. Breuer G, de Jaeger L, Artus VPG, Martens DE, Springer J, Draaisma RB, et al. Superior triacylglycerol (TAG) accumulation in starchless mutants of *Scenedesmus obliquus*: (II) evaluation of TAG yield and productivity in controlled photobioreactors. *Biotechnol Biofuels*. 2014;7:70.
56. Bogen C, Klassen V, Wichmann J, La Russa M, Doebbe A, Grundmann M, et al. Identification of *Monoraphidium contortum* as a promising species for liquid biofuel production. *Bioresour Technol*. 2013;133:622–6.
57. Gouveia JD, Ruiz J, van den Broek LAM, Hesselink T, Peters S, Kleinegris DMM, et al. *Botryococcus braunii* strains compared for biomass productivity, hydrocarbon and carbohydrate content. *J Biotechnol*. 2017;248:77–86.
58. Benvenuti G, Ruiz J, Lamers PP, Bosma R, Wijffels RH, Barbosa MJ. Towards microalgal triglycerides in the commodity markets. *Biotechnol Biofuels*. 2017;10:188.
59. Gonzalez-Fernandez C, Sialve B, Molinuevo-Salces B. Anaerobic digestion of microalgal biomass: challenges, opportunities and research needs. *Bioresour Technol*. 2015;198:896–906.
60. Santos-Ballardo DU, Rossi S, Reyes-Moreno C, Valdez-Ortiz A. Microalgae potential as a biogas source: current status, restraints and future trends. *Rev Environ Sci Biotechnol*. 2016;15:243–64.
61. Mahdy A, Mendez L, Blanco S, Ballesteros M, González-Fernández C. Protease cell wall degradation of *Chlorella vulgaris*: effect on methane production. *Bioresour Technol*. 2014;171:421–7.
62. Mahdy A, Mendez L, Tomás-Pejó E, del Mar Morales M, Ballesteros M, González-Fernández C. Influence of enzymatic hydrolysis on the biochemical methane potential of *Chlorella vulgaris* and *Scenedesmus* sp. *J Chem Technol Biotechnol*. 2016;91:1299–305.
63. Sharma A, Arya SK. Hydrogen from algal biomass: a review of production process. *Biotechnol Rep*. 2017;15:63–9.
64. Khetkorn W, Rastogi RP, Incharoensakdi A, Lindblad P, Madamwar D, Pandey A, et al. Microalgal hydrogen production—a review. *Bioresour Technol*. 2017;243:1194–206.
65. Dubini A, Ghirardi ML. Engineering photosynthetic organisms for the production of biohydrogen. *Photosynth Res*. 2015;123:241–53.
66. Venkanna D, Sudfeld C, Baier T, Homburg SV, Patel AV, Wobbe L, et al. Knock-down of the IFR1 protein perturbs the homeostasis of reactive electrophile species and boosts photosynthetic hydrogen production in *Chlamydomonas reinhardtii*. *Front Plant Sci*. 2017;8:1347.
67. Bayro-Kaiser V, Nelson N. Temperature-sensitive PSII: a novel approach for sustained photosynthetic hydrogen production. *Photosynth Res*. 2016;130:113–21.
68. Cheah WY, Ling TC, Juan JC, Lee DJ, Chang JS, Show PL. Biorefineries of carbon dioxide: from carbon capture and storage (CCS) to bioenergies production. *Bioresour Technol*. 2016;215:346–56.
69. da Rosa GM, Moraes L, Cardias BB, Souza da RAZ de M, Costa JAV. Chemical absorption and CO₂ biofixation via the cultivation of *Spirulina* in semicontinuous mode with nutrient recycle. *Bioresour Technol*. 2015;192:321–7.
70. Assunção J, Batista AP, Manoel J, da Silva TL, Marques P, Reis A, et al. CO₂ utilization in the production of biomass and biocompounds by three different microalgae. *Eng Life Sci*. 2017;17:1126–35.
71. Adamczyk M, Lasek J, Skawińska A. CO₂ biofixation and growth kinetics of *Chlorella vulgaris* and *Nannochloropsis gaditana*. *Appl Biochem Biotechnol*. 2016;179:1248–61.
72. Caporgno MP, Taleb A, Olkiewicz M, Font J, Pruvost J, Legrand J, et al. Microalgae cultivation in urban wastewater: nutrient removal and biomass production for biodiesel and methane. *Algal Res*. Elsevier. 2015;10:232–9.
73. Lv J, Feng J, Liu Q, Xie S. Microalgal cultivation in secondary effluent: recent developments and future work. *Int J Mol Sci*. 2017;18:79.
74. Leu S, Boussiba S. Advances in the production of high-value products by microalgae. *Ind Biotechnol*. 2014;10:169–83.
75. Chew KW, Yap JY, Show PL, Suan NH, Juan JC, Ling TC, et al. Microalgal biorefinery: high value products perspectives. *Bioresour Technol*. Elsevier. 2017;229:53–62.
76. 't Lam GP, Vermuë MH, Eppink MHM, Wijffels RH, van den Berg C. Multi-product microalgae biorefineries: from concept towards reality. *Trends Biotechnol*. 2018;36:216–27.
77. Fernandes BD, Mota A, Teixeira JA, Vicente AA. Continuous cultivation of photosynthetic microorganisms: approaches, applications and future trends. *Biotechnol Adv*. 2015;33:1228–45.

78. Cuaresma M, Janssen M, Vilchez C, Wijffels RH. Horizontal or vertical photobioreactors? How to improve microalgae photosynthetic efficiency. *Bioresour Technol*. 2011;110:5129–37.
79. Münkler R, Schmid-Staiger U, Werner A, Hirth T. Optimization of outdoor cultivation in flat panel airlift reactors for lipid production by *Chlorella vulgaris*. *Biotechnol Bioeng*. 2013;110:2882–93.
80. Wang B, Lan CQ, Horsman M. Closed photobioreactors for production of microalgal biomasses. *Biotechnol Adv*. 2012;30:904–12.
81. Ozkan A, Kinney K, Katz L, Berberoglu H. Reduction of water and energy requirement of algae cultivation using an algae biofilm photobioreactor. *Bioresour Technol*. 2012;114:542–8.
82. Zhang Q, Liu C, Li Y, Yu Z, Chen Z, Ye T, et al. Cultivation of algal biofilm using different lignocellulosic materials as carriers. *Biotechnol Biofuels*. 2017;10:1–16.
83. Wrede D, Taha M, Miranda AF, Kadali K, Stevenson T, Ball AS, et al. Co-cultivation of fungal and microalgal cells as an efficient system for harvesting microalgal cells, lipid production and wastewater treatment. *PLoS ONE*. 2014;9:e113497.
84. Chen CL, Chang JS, Lee DJ. Dewatering and drying methods for microalgae. *Dry Technol*. 2015;33:443–54.
85. Yaakob Z, Ali E, Zainal A, Mohamad M, Takriff MS. An overview: biomolecules from microalgae for animal feed and aquaculture. *J Biol Res*. 2014;21:6.
86. Bonente G, Passarini F, Cazzaniga S, Mancone C, Buia MC, Tripodi M, et al. The occurrence of the *psbS* gene product in *Chlamydomonas reinhardtii* and in other photosynthetic organisms and its correlation with energy quenching. *Photochem Photobiol*. 2008;84:1359–70.
87. Peers G, Truong TB, Ostendorf E, Busch A, Elrad D, Grossman AR, et al. An ancient light-harvesting protein is critical for the regulation of algal photosynthesis. *Nature*. 2009;462:518–21.
88. Kulheim C, Agren J, Jansson S. Rapid regulation of light harvesting and plant fitness in the field. *Science*. 2002;297:91–3.
89. Formighieri C, Franck F, Bassi R. Regulation of the pigment optical density of an algal cell: filling the gap between photosynthetic productivity in the laboratory and in mass culture. *J Biotechnol*. 2012;162:115–23.
90. Whitney SM, Houtz RL, Alonso H. Advancing our understanding and capacity to engineer nature's CO₂-sequestering enzyme, Rubisco. *Plant Physiol*. 2011;155:27–35.
91. Reinfelder JR. Carbon concentrating mechanisms in eukaryotic marine phytoplankton. *Ann Rev Mar Sci*. 2011;3:291–315.
92. Langridge P. Reinventing the green revolution by harnessing crop mutant resources. *Plant Physiol*. 2014;166:1682–3.
93. Gaut BS, Seymour DK, Liu Q, Zhou Y. Demography and its effects on genomic variation in crop domestication. *Nat Plants*. 2018;4:512–20.
94. Snow AA, Smith VH. Genetically engineered algae for biofuels: a key role for ecologists. *Bioscience*. 2012;62:765–8.
95. Nakajima Y, Ueda R. Improvement of photosynthesis in dense microalgal suspension by reduction of light harvesting pigments. *J Appl Phycol*. 1997;9:503–10.
96. Beckmann J, Lehr F, Finazzi G, Hankamer B, Posten C, Wobbe L, et al. Improvement of light to biomass conversion by de-regulation of light-harvesting protein translation in *Chlamydomonas reinhardtii*. *J Biotechnol*. 2009;142:70–7.
97. Bonente G, Formighieri C, Mantelli M, Catalanotti C, Giuliano G, Morosinotto T, et al. Mutagenesis and phenotypic selection as a strategy toward domestication of *Chlamydomonas reinhardtii* strains for improved performance in photobioreactors. *Photosynth Res*. 2011;108:107–20.
98. Perrine Z, Negi S, Sayre RT. Optimization of photosynthetic light energy utilization by microalgae. *Algal Res*. 2012;1:134–42.
99. Kirst H, Garcia-Cerdan JG, Zurbriggen A, Ruehle T, Melis A. Truncated photosystem chlorophyll antenna size in the green microalga *Chlamydomonas reinhardtii* upon deletion of the *TLA3-CpSRP43* gene. *Plant Physiol*. 2012;160:2251–60.
100. Jeong J, Baek K, Kirst H, Melis A, Jin E. Loss of CpSRP54 function leads to a truncated light-harvesting antenna size in *Chlamydomonas reinhardtii*. *Biochim Biophys Acta Bioenerg*. 2017;1858:45–55.
101. Cazzaniga S, Dall'Osto L, Szaub J, Scibilia L, Ballottari M, Purton S, et al. Domestication of the green alga *Chlorella sorokiniana*: reduction of antenna size improves light-use efficiency in a photobioreactor. *Biotechnol Biofuels*. 2014;7:157.
102. Perin G, Bellan A, Segalla A, Meneghesso A, Alboresi A, Morosinotto T. Generation of random mutants to improve light-use efficiency of *Nannochloropsis gaditana* cultures for biofuel production. *Biotechnol Biofuels*. 2015;8:161.
103. Oey M, Ross IL, Stephens E, Steinbeck J, Wolf J, Radzun KA, et al. RNAi Knock-Down of LHCBM1, 2 and 3 Increases Photosynthetic H₂ Production Efficiency of the Green Alga *Chlamydomonas reinhardtii*. *PLoS ONE*. 2013;8:e61375.
104. Ho M-Y, Shen G, Canniffe DP, Zhao C, Bryant DA. Light-dependent chlorophyll f synthase is a highly divergent paralog of PsbA of photosystem II. *Science*. 2016;353:aaf9178.
105. Tchernov D, Livne A, Kaplan A, Sukenik A. The kinetic properties of ribulose-1,5-bisphosphate carboxylase/oxygenase may explain the high apparent photosynthetic affinity of *Nannochloropsis* sp. to ambient inorganic carbon. *Isr J Plant Sci*. 2008;56:37–44.
106. John Andrews T, Whitney SM. Manipulating ribulose bisphosphate carboxylase/oxygenase in the chloroplasts of higher plants. *Arch Biochem Biophys*. 2003;414:159–69.
107. Wilson RH, Martin-Avila E, Conlan C, Whitney SM. An improved *Escherichia coli* screen for Rubisco identifies a protein-protein interface that can enhance CO₂-fixation kinetics. *J Biol Chem*. 2018;293:18–27.
108. Tsai Y-CC, Lapina MC, Bhushan S, Mueller-Cajar O. Identification and characterization of multiple rubisco activases in chemoautotrophic bacteria. *Nat Commun*. 2015;6:8883.
109. Wakabayashi K, Misawa Y, Mochiji S, Kamiya R. Reduction-oxidation poise regulates the sign of phototaxis in *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci USA*. 2011;108:11280–4.
110. Kim JYH, Kwak HS, Sung YJ, Choi HI, Hong ME, Lim HS, et al. Microfluidic high-throughput selection of microalgal strains with superior photosynthetic productivity using competitive phototaxis. *Sci Rep*. 2016;6:21155.
111. Kromdijk J, Glowacka K, Leonelli L, Gabilly ST, Iwai M, Niyogi KK, et al. Improving photosynthesis and crop productivity by accelerating recovery from photoprotection. *Science*. 2016;354:857–61.
112. Glowacka K, Kromdijk J, Kucera K, Xie J, Cavanagh AP, Leonelli L, et al. Photosystem II subunit S overexpression increases the efficiency of water use in a field-grown crop. *Nat Commun*. 2018;9:868.
113. Peers G. Enhancement of biomass production by disruption of light energy dissipation pathways, U.S. Patent No. 8,940,508. Washington, DC: Patent and Trademark Office; 2015. <https://app.dimensions.ai/details/patent/US-8940508-B2>.
114. Truong A. Investigating the role(s) of LHCSR in *Chlamydomonas reinhardtii*. Ph.D. thesis. University of California, Berkeley.
115. Berteotti S, Ballottari M, Bassi R. Increased biomass productivity in green algae by tuning non-photochemical quenching. *Sci Rep*. 2016;6:21339.
116. Cantrell M, Peers G. A mutant of *Chlamydomonas* without LHCSR maintains high rates of photosynthesis, but has reduced cell division rates in sinusoidal light conditions. *PLoS ONE*. 2017;12:e0179395.
117. Förster B, Osmond CB, Pogson BJ. Improved survival of very high light and oxidative stress is conferred by spontaneous gain-of-function mutations in *Chlamydomonas*. *Biochim Biophys Acta Bioenerg*. 2005;1709:45–57.
118. Fischer BB, Ledford HK, Wakao S, Huang SG, Casero D, Pellegrini M, et al. Singlet oxygen resistant 1 links reactive electrophile signaling to singlet oxygen acclimation in *Chlamydomonas reinhardtii*. *Proc Natl Acad Sci*. 2012;109:1302–11.
119. Schierenbeck L, Ries D, Rogge K, Grewe S, Weisshaar B, Kruse O. Fast forward genetics to identify mutations causing a high light tolerant phenotype in *Chlamydomonas reinhardtii* by whole-genome-sequencing. *BMC Genomics*. 2015;16:57.
120. Kang S, Kim K-H, Kim Y-C. A novel electroporation system for efficient molecular delivery into *Chlamydomonas reinhardtii* with a 3-dimensional microelectrode. *Sci Rep*. 2015;5:15835.
121. Mini P, Demurtas OC, Valentini S, Pallara P, Aprea G, Ferrante P, et al. Agrobacterium-mediated and electroporation-mediated transformation of *Chlamydomonas reinhardtii*: a comparative study. *BMC Biotechnol*. 2018;18:11.
122. Dunahay TG. Transformation of *Chlamydomonas reinhardtii* with silicon carbide whiskers. *Biotechniques*. 1993;15:452–5, 457–8, 460.

123. Kim S, Lee Y-C, Cho D-H, Lee HU, Huh YS, Kim G-J, et al. A simple and non-invasive method for nuclear transformation of intact-walled *Chlamydomonas reinhardtii*. *PLoS ONE*. 2014;9:e101018.
124. Somchai P, Jitrakorn S, Thitamadee S, Meetam M, Saksmerprome V. Use of microalgae *Chlamydomonas reinhardtii* for production of double-stranded RNA against shrimp virus. *Aquac Rep*. 2016;3:178–83.
125. Kumar M, Jeon J, Choi J, Kim S-R. Rapid and efficient genetic transformation of the green microalga *Chlorella vulgaris*. *J Appl Phycol*. 2018;30:1735–45.
126. Chungjatupornchai W, Kitraksa P, Fa-aroonsawat S. Stable nuclear transformation of the oleaginous microalga *Neochloris oleoabundans* by electroporation. *J Appl Phycol*. 2016;28:191–9.
127. Kathiresan S, Chandrashekar A, Ravishankar GA, Sarada R. Regulation of astaxanthin and its intermediates through cloning and genetic transformation of β -carotene ketolase in *Haematococcus pluvialis*. *J Biotechnol*. 2015;196–197:33–41.
128. Kindle KL. Nuclear transformation: technology and applications. In: Rochaix JD, Goldschmidt-Clermont M, Merchant S, editors. The molecular biology of chloroplasts mitochondria in *Chlamydomonas*. Dordrecht: Springer; 1998. p. 41–61.
129. Schroda M. RNA silencing in *Chlamydomonas*: mechanisms and tools. *Curr Genet*. 2006;49:69–84.
130. Zhang H, Lang Z, Zhu J-K. Dynamics and function of DNA methylation in plants. *Nat Rev Mol Cell Biol*. 2018;19:489–506.
131. Kim MY, Zilberman D. DNA methylation as a system of plant genomic immunity. *Trends Plant Sci*. 2014;19:320–6.
132. Rasala BA, Lee PA, Shen Z, Briggs SP, Mendez M, Mayfield SP. Robust expression and secretion of Xylanase1 in *Chlamydomonas reinhardtii* by fusion to a selection gene and processing with the FMDV 2A peptide. *PLoS ONE*. 2012;7:e43349.
133. Onishi M, Pringle JR. Robust transgene expression from bicistronic mRNA in the green alga *Chlamydomonas reinhardtii*. G3 (Bethesda). 2016;6:4115–25.
134. Weiner I, Atar S, Schweitzer S, Eilenberg H, Feldman Y, Avitan M, et al. Enhancing heterologous expression in *Chlamydomonas reinhardtii* by transcript sequence optimization. *Plant J*. 2018;94:22–31.
135. Kim JH, Lee S-R, Li L-H, Park H-J, Park J-H, Lee KY, et al. High cleavage efficiency of a 2A peptide derived from porcine teschovirus-1 in human cell lines, Zebrafish and Mice. *PLoS ONE*. 2011;6:e18556.
136. Kwon S, Kang NK, Koh HG, Shin S-E, Lee B, Jeong B, et al. Enhancement of biomass and lipid productivity by overexpression of a bZIP transcription factor in *Nannochloropsis salina*. *Biotechnol Bioeng*. 2018;115:331–40.
137. Doron L, Segal N, Shapira M. Transgene expression in microalgae—from tools to applications. *Front Plant Sci*. 2016;7:505.
138. Wannathong T, Waterhouse JC, Young REB, Economou CK, Purton S. New tools for chloroplast genetic engineering allow the synthesis of human growth hormone in the green alga *Chlamydomonas reinhardtii*. *Appl Microbiol Biotechnol*. 2016;100:5467–77.
139. Nymark M, Sharma AK, Sparstad T, Bones AM, Winge P. A CRISPR/Cas9 system adapted for gene editing in marine algae. *Sci Rep*. 2016;6:24951.
140. Baek K, Kim DH, Jeong J, Sim SJ, Melis A, Kim J-S, et al. DNA-free two-gene knockout in *Chlamydomonas reinhardtii* via CRISPR-Cas9 ribonucleoproteins. *Sci Rep*. 2016;6:30620.
141. Shin SE, Lim JM, Koh HG, Kim EK, Kang NK, Jeon S, et al. CRISPR/Cas9-induced knockout and knock-in mutations in *Chlamydomonas reinhardtii*. *Sci Rep*. 2016;6:27810.
142. Ferenczi A, Pyott DE, Xipnitou A, Molnar A. Efficient targeted DNA editing and replacement in *Chlamydomonas reinhardtii* using Cpf1 ribonucleoproteins and single-stranded DNA. *Proc Natl Acad Sci USA*. 2017;114:13567–72.
143. Wang Q, Lu Y, Xin Y, Wei L, Huang S, Xu J. Genome editing of model oleaginous microalgae *Nannochloropsis* spp. by CRISPR/Cas9. *Plant J*. 2016;88:1071–81.
144. Li F, Gao D, Hu H. High-efficiency nuclear transformation of the oleaginous marine *Nannochloropsis* species using PCR product. *Biosci Biotechnol Biochem*. 2014;78:812–7.
145. Ajjawi I, Verruto J, Aquí M, Soriaga LB, Coppersmith J, Kwok K, et al. Lipid production in *Nannochloropsis gaditana* is doubled by decreasing expression of a single transcriptional regulator. *Nat Biotechnol*. 2017;35:647–52.
146. Mitra M, Kirst H, Dewez D, Melis A. Modulation of the light-harvesting chlorophyll antenna size in *Chlamydomonas reinhardtii* by TLA1 gene over-expression and RNA interference. *Philos Trans R Soc B Biol Sci*. 2012;367:3430–43.
147. Berger H, de Mia M, Morisse S, Marchand C, Lemaire SD, Wobbe L, et al. A light switch based on protein S-nitrosylation fine-tunes photosynthetic light-harvesting in the microalga *Chlamydomonas reinhardtii*. *Plant Physiol*. 2016;171:01878.
148. Ducat DC, Silver PA. Improving carbon fixation pathways. *Curr Opin Chem Biol*. 2012;16:337–44.
149. Carmo-Silva E, Scales JC, Madgwick PJ, Parry MAJ. Optimizing Rubisco and its regulation for greater resource use efficiency. *Plant Cell Environ*. 2015;38:1817–32.
150. Larson EM, O'Brien CM, Zhu G, Spreitzer RJ, Portis AR. Specificity for activase is changed by a Pro-89 to Arg substitution in the large subunit of ribulose-1,5-bisphosphate carboxylase/oxygenase. *J Biol Chem*. 1997;272:17033–7.
151. Ott CM, Smith BD, Portis AR, Spreitzer RJ. Activase region on chloroplast ribulose-1,5-bisphosphate carboxylase/oxygenase. *J Biol Chem*. 2000;275:26241–4.
152. Li C, Salvucci ME, Portis AR. Two residues of rubisco activase involved in recognition of the rubisco substrate. *J Biol Chem*. 2005;280:24864–9.
153. Wei L, Wang Q, Xin Y, Lu Y, Xu J. Enhancing photosynthetic biomass productivity of industrial oleaginous microalgae by overexpression of RuBisCO activase. *Algal Res*. 2017;27:366–75.
154. Pinto TS, Malcata FX, Arrabaça JD, Silva JM, Spreitzer RJ, Esquivel MG. Rubisco mutants of *Chlamydomonas reinhardtii* enhance photosynthetic hydrogen production. *Appl Microbiol Biotechnol*. 2013;97:5635–43.
155. Zhu G, Kurek I, Liu L. Engineering photosynthetic enzymes involved in CO₂-assimilation by gene shuffling. In: Rebeiz CA, Benning C, Bohnert HJ, Daniell H, Hooper JK, Lichtenthaler HK, Portis AR Jr, Tripathy BC, editors. The chloroplast: basics and application (advances in photosynthesis and respiration), Chap. 20. Dordrecht: Springer; 2010. p. 307–22.
156. Johnson X, Wostrickoff K, Finazzi G, Kuras R, Schwarz C, Bujaldon S, et al. MRL1, a conserved pentatricopeptide repeat protein, is required for stabilization of rbcL mRNA in *Chlamydomonas* and *Arabidopsis*. *Plant Cell*. 2010;22:234–48.
157. Fang L, Lin HX, Low CS, Wu MH, Chow Y, Lee YK. Expression of the *Chlamydomonas reinhardtii* Sedoheptulose-1,7-bisphosphatase in *Dunaliella bardawil* leads to enhanced photosynthesis and increased glycerol production. *Plant Biotechnol J*. 2012;10:1129–35.
158. Spalding MH. Modulation of low carbon dioxide inducible proteins (lci) for increased biomass production and photosynthesis. U.S. patent 2012. <https://patents.google.com/patent/WO2013006361A1>.
159. Costas AMG, White AK, Metcalf WW. Purification and characterization of a novel phosphorus-oxidizing enzyme from *Pseudomonas stutzeri* WM88. *J Biol Chem*. 2001;276:17429–36.
160. Loera-Quezada MM, Leyva-González MA, Velázquez-Juárez G, Sanchez-Calderón L, Do Nascimento M, López-Arredondo D, et al. A novel genetic engineering platform for the effective management of biological contaminants for the production of microalgae. *Plant Biotechnol J*. 2016;14:2066–76.
161. Doebe A, Rupprecht J, Beckmann J, Mussnug JH, Hallmann A, Hankamer B, et al. Functional integration of the HUP1 hexose symporter gene into the genome of *C. reinhardtii*: impacts on biological H₂ production. *J Biotechnol*. 2007;131:27–33.
162. Leite GB, Paranjape K, Abdelaziz AEM, Hallenbeck PC. Utilization of biodiesel-derived glycerol or xylose for increased growth and lipid production by indigenous microalgae. *Bioresour Technol*. 2015;184:123–30.
163. Ramos-Martinez EM, Fimognari L, Sakuragi Y. High-yield secretion of recombinant proteins from the microalga *Chlamydomonas reinhardtii*. *Plant Biotechnol J*. 2017;15:1214–24.
164. Imam SH, Buchanan MJ, Shin HC, Snell WJ. The *Chlamydomonas* cell wall: characterization of the wall framework. *J Cell Biol*. 1985;101:1599–607.
165. Malinovsky FG, Fangel JU, Willats WGT. The role of the cell wall in plant immunity. *Front Plant Sci*. 2014;5:178.
166. Benedetti M, Pontiggia D, Raggi S, Cheng Z, Scaloni F, Ferrari S, et al. Plant immunity triggered by engineered in vivo release of

- oligogalacturonides, damage-associated molecular patterns. *Proc Natl Acad Sci*. 2015;112:5533–8.
167. Blifernez-Klassen O, Klassen V, Doebbe A, Kersting K, Grimm P, Wobbe L, et al. Cellulose degradation and assimilation by the unicellular phototrophic eukaryote *Chlamydomonas reinhardtii*. *Nat Commun*. 2012;3:1214.
168. Kricka W, Fitzpatrick J, Bond U. Metabolic engineering of yeasts by heterologous enzyme production for degradation of cellulose and hemicellulose from biomass: a perspective. *Front Microbiol*. 2014;5:174.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

