



Local boundedness for solutions of a class of nonlinear elliptic systems

Giovanni Cupini¹ · Francesco Leonetti² · Elvira Mascolo³

Received: 14 October 2021 / Accepted: 21 February 2022
© The Author(s) 2022

Abstract

In this paper we are concerned with the regularity of solutions to a nonlinear elliptic system of m equations in divergence form, satisfying p growth from below and q growth from above, with $p \leq q$; this case is known as p, q -growth conditions. Well known counterexamples, even in the simpler case $p = q$, show that solutions to systems may be singular; so, it is necessary to add suitable structure conditions on the system that force solutions to be regular. Here we obtain local boundedness of solutions under a componentwise coercivity condition. Our result is obtained by proving that each component u^α of the solution $u = (u^1, \dots, u^m)$ satisfies an improved Caccioppoli's inequality and we get the boundedness of u^α by applying De Giorgi's iteration method, provided the two exponents p and q are not too far apart. Let us remark that, in dimension $n = 3$ and when $p = q$, our result works for $\frac{3}{2} < p \leq 3$, thus it complements the one of Bjorn whose technique allowed her to deal with $p \leq 2$ only. In the final section, we provide applications of our result.

Mathematics Subject Classification Primary: 35J47 · Secondary: 35B65

Communicated by Y. Giga.

✉ Francesco Leonetti
francesco.leonetti@univaq.it

Giovanni Cupini
giovanni.cupini@unibo.it

Elvira Mascolo
elvira.mascolo@unifi.it

¹ Dipartimento di Matematica, Università di Bologna, Piazza di Porta S. Donato 5, 40126 Bologna, Italy

² Dipartimento di Ingegneria e Scienze dell'Informazione e Matematica, Università di L'Aquila, Via Vetoio snc - Coppito, 67100 L'Aquila, Italy

³ Dipartimento di Matematica e Informatica "U. Dini", Università di Firenze, Viale Morgagni 67/A, 50134 Firenze, Italy

1 Introduction

In this paper we are concerned with the regularity of solutions to a nonlinear elliptic system of m equations in divergence form

$$\sum_{i=1}^n \frac{\partial}{\partial x_i} (A_i^\alpha(x, Du(x))) = 0, \quad 1 \leq \alpha \leq m, \tag{1.1}$$

where $x \in \Omega$ and Ω is a bounded open set in \mathbb{R}^n , $n \geq 2$. The function $u : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}^m$, has components (u^1, \dots, u^m) ; then $Du(x)$ is the $m \times n$ matrix $\left(\frac{\partial u^\alpha}{\partial x_i}(x) \right)_{\substack{\alpha=1, \dots, m \\ i=1, \dots, n}}$.

We assume that $A_i^\alpha : \Omega \times \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$, $1 \leq i \leq n$, $1 \leq \alpha \leq m$, are Carathéodory functions satisfying for every $x \in \Omega$ and for every $z = (z^1, \dots, z^m)^T \in \mathbb{R}^{m \times n}$ the following p, q -growth assumptions:

$$v|z^\alpha|^p - a(x) \leq \sum_{i=1}^n A_i^\alpha(x, z)z_i^\alpha \quad \forall \alpha \in \{1, \dots, m\}, \tag{1.2}$$

$$\sum_{i=1}^n |A_i^\alpha(x, z)| \leq M(|z|^{q-1} + b(x)), \tag{1.3}$$

where $1 < p \leq q$, $p \leq n$, $v, M > 0$, $a \in L_{loc}^{\tau_1}(\Omega)$ and $b \in L_{loc}^{\tau_2}(\Omega)$ are non-negative functions, with $1 < \tau_i \leq +\infty$, $i = 1, 2$, and $\tau_2 \geq \frac{q}{q-1}$.

Let us recall that $u \in W_{loc}^{1,q}(\Omega; \mathbb{R}^m)$ is a weak solution of (1.1) if

$$\int_B \sum_{\beta=1}^m \sum_{i=1}^n A_i^\beta(x, Du(x)) D_i \psi^\beta(x) dx = 0, \tag{1.4}$$

for every open set $B \Subset \Omega$ and for every $\psi \in W_0^{1,q}(B; \mathbb{R}^m)$.

As usual the Sobolev exponent is $p^* = \frac{np}{n-p}$ if $p < n$, and p^* is any real number $\mu > p$ if $p = n$. The Hölder conjugate exponent of p is $p' = \frac{p}{p-1}$. We use the position $\frac{1}{+\infty} = 0$. Our regularity result is the following.

Theorem 1.1 *Assume that (1.2) and (1.3) hold, with $1 < p \leq n$, $p \leq q$ and $1 < \tau_1, \tau_2 \leq +\infty$, satisfying*

$$q < p^* \frac{n}{p(n+1)}, \quad \tau_1 > \frac{n}{p}, \quad \tau_2 \geq \frac{q}{q-1}. \tag{1.5}$$

Then any weak solution $u \in W_{loc}^{1,q}(\Omega; \mathbb{R}^m)$ of (1.1) is locally bounded.

In the vector-valued case, as suggested by well known counterexamples [11, 13, 14, 18–20, 23, 24, 28, 38, 43, 45, 49, 50], special structures on the operator are required for everywhere regularity, even under reasonable assumptions on the coefficients; see also the surveys [40, 41] and [25].

In the literature there are still few contributions about the boundedness of weak solutions to elliptic systems. Ladyzhenskaya and Ural'tseva ([27], Chapter 7) first proposed the local

boundedness of solutions $u = (u^1, u^2, \dots, u^m)$ to the *linear* elliptic system

$$\sum_{i=1}^n \frac{\partial}{\partial x_i} \left(\sum_{j=1}^n a_{ij}(x) u_{x_j}^\alpha + \sum_{\beta=1}^m b_i^{\alpha\beta}(x) u^\beta + f_i^\alpha(x) \right) + \sum_{i=1}^n \sum_{\beta=1}^m c_i^{\alpha\beta}(x) u_{x_i}^\beta + \sum_{\beta=1}^m d^{\alpha\beta}(x) u^\beta = f^\alpha(x), \quad \forall \alpha = 1, 2, \dots, m, \tag{1.6}$$

with bounded measurable coefficients $a_{ij}, b_i^{\alpha\beta}, c_i^{\alpha\beta}, d^{\alpha\beta}$ and given functions f_i^α, f^α . Here the *structure condition* is stated in terms of the positive definite $n \times n$ matrix (a_{ij}) , which *does not depend* on α, β . In [39] Meier extended these results to nonlinear elliptic systems of the form

$$\sum_{i=1}^n \frac{\partial}{\partial x_i} (A_i^\alpha(x, u, Du)) = 0, \tag{1.7}$$

under the following p -growth conditions, $1 < p \leq n$,

$$\sum_{i=1}^n \sum_{\alpha=1}^m A_i^\alpha(x, u, z) z_i^\alpha \geq |z|^p - d(x)|u|^p - g(x) \tag{1.8}$$

$$|A^\alpha(x, u, z)| \leq a|z|^{p-1} + b(x)|u|^{p-1} + e(x) \tag{1.9}$$

for $a > 0$ and under suitable integrability assumptions on the nonnegative functions b, e, d, g . Meier introduces the so-called *indicator function* of the operator

$$I_A(x, u, Du) := \sum_{\alpha, \beta, i} A_i^\alpha(x, u, Du) D_i u^\beta \frac{u^\alpha u^\beta}{|u|^2} \tag{1.10}$$

and a pointwise assumption turns out to be crucial in Meier’s techniques, indeed a weak solution u of (1.7) is locally bounded if

$$I_A(x, u, Du) \geq 0 \tag{1.11}$$

holds for large values of $|u|$. Notice that (1.11) is satisfied in linear case (1.6). Assumption (1.11) is satisfied also by some nonlinear operators. For example:

$$A_i^\alpha(Du) = \sigma(Du) D_i u^\alpha, \tag{1.12}$$

when $0 \leq \sigma$, like in the case of Euler’s system of the functional

$$\int F(|Du|) dx, \tag{1.13}$$

where F increases and we take $\sigma(Du) = \frac{F'(|Du|)}{|Du|}$. A third example, for which (1.11) holds true, is given when considering Euler’s system of the anisotropic integral

$$\int \sum_{i=1}^n g_i(|D_i u|) dx, \tag{1.14}$$

where g_i increases and we take

$$A_i^\alpha(Du) = \frac{g_i'(|D_i u|)}{|D_i u|} D_i u^\alpha, \tag{1.15}$$

see section 4 in [30]. Let us look at another example: we set $m = n$ and we consider the polyconvex integral

$$\int (|Du|^p + h(\det Du))dx, \tag{1.16}$$

where h is convex, C^1 , bounded from below. In this case Euler’s system gives

$$A_i^\alpha(Du) = p|Du|^{p-2}D_iu^\alpha + h'(\det Du)(\text{Cof } Du)_i^\alpha, \tag{1.17}$$

where $(\text{Cof } Du)_i^\alpha$ is the determinant of the $(n - 1) \times (n - 1)$ matrix obtained from the $n \times n$ matrix Du by deleting row α and column i , with the sign given by $(-1)^{\alpha+i}$. It turns out that

$$I_A(x, u, Du) = p|Du|^{p-2} \sum_{i=1}^n \left(\sum_{\alpha=1}^n \frac{u^\alpha}{|u|} D_iu^\alpha \right)^2 + h'(\det Du) \det Du \geq \inf_{\mathbb{R}} h - h(0), \tag{1.18}$$

then we get (1.11), provided $h(0) = \inf_{\mathbb{R}} h$; see Sect. 3, later in the present paper; see also [31].

The previous examples show that Meier’s condition allows us to deal with quite a large class of nonlinear systems. Boundedness results for weak solutions to nonlinear elliptic systems are proved by Krömer [26] under assumptions similar to (1.11), see also Landes [29].

Actually Meier’s regularity result is obtained under a weaker assumption, since I_A can be allowed to be negative, but not too much.

More precisely, under (1.7) and (1.8), there exist positive constants λ and L such that

$$I_A(x, u, z) := \sum_{\alpha,\beta,i} A_i^\alpha(x, u, z) z_i^\beta \frac{u^\alpha u^\beta}{|u|^2} \geq - \left\{ \delta |z|^p + \left(\frac{1}{\delta} \right)^\lambda [d(x)|u|^p + g(x)] \right\} \tag{1.19}$$

for every $\delta \in (0, 1)$, for all $(x, u, z) \in \Omega \times \mathbb{R}^m \times \mathbb{R}^{m \times n}$, with $|u| > L$.

Let us observe that the following linear decoupled system does not verify (1.19), see [32] and Sect. 3, later in the present paper:

$$A_i^\alpha(x, Du) = \sigma^\alpha(x) D_iu^\alpha, \tag{1.20}$$

where $m = 2$,

$$\sigma^1(x) = 18 + 2 \sin(|x|^2) \quad \text{and} \quad \sigma^2(x) = 2 + \sin(|x|^2). \tag{1.21}$$

Now we consider another example, see [31], in which the equations are coupled and Meier’s condition (1.19) is not satisfied: it is Euler’s system of

$$\int [|Du|^2 + h(D_1u^1 D_1u^2)] dx \tag{1.22}$$

where $m = 2$, h is convex, C^1 , bounded from below, so that

$$A_i^\alpha(Du) = 2D_iu^\alpha + h'(D_1u^1 D_1u^2) D_1u^{\hat{\alpha}} \delta_{i1}, \tag{1.23}$$

where

$$\hat{\alpha} = 2 \text{ if } \alpha = 1 \text{ and } \hat{\alpha} = 1 \text{ if } \alpha = 2; \text{ moreover, } \delta_{i1} = 1 \text{ if } i = 1 \text{ and } \delta_{i1} = 0 \text{ otherwise.}$$

Meier’s condition (1.19) is not satisfied, provided $h'(0) \leq -8$: for instance, $h(t) = 16\sqrt{1 + (t - 1)^2}$; see Sect. 3 later in the present paper.

Combining coefficients $\sigma^\alpha(x)$ similar to (1.21) with the nonlinear part of (1.23), we are able to build an example with p growth that does not satisfy Meier’s condition (1.19). Indeed,

$$A_i^\alpha(Du) = \sigma^\alpha(x)p|Du|^{p-2}D_iu^\alpha + h'(D_1u^1D_1u^2)D_1u^{\hat{\alpha}}\delta_{i1}, \tag{1.24}$$

where $2 \leq p, h$ is convex, C^1 , bounded from below; $\hat{\alpha}$ and δ_{i1} are defined as before. Moreover, $m = 2$ and

$$\sigma^1(x) = 48 + 3 \sin(|x|^2) \text{ and } \sigma^2(x) = 2 + \sin(|x|^2). \tag{1.25}$$

Meier’s condition (1.19) is not satisfied, provided $h'(0) \leq 0$: for instance, $h(t) = (1 + t^2)^{p/4}$; see Sect. 3 for the details.

In [2] Bjorn obtained boundedness of solutions u of systems without considering the indicator function but assuming componentwise coercivity:

$$\nu|z^\alpha|^p - a(x) - b(x)|u|^p \leq \sum_{i=1}^n A_i^\alpha(x, u, z)z_i^\alpha, \text{ with } \nu > 0. \tag{1.26}$$

Previous assumption (1.26) says that, even if row α of the system contains all the components of $z = Du$, after multiplying this row by component α of $z = Du$, from below we only see the α component of $z = Du$ and none of other components.

(1.26) is satisfied in system (1.20), provided $\sigma^\alpha(x) \geq \nu$ for some positive constant ν . Furthermore, the structure in (1.12) guarantees (1.26), provided $\sigma(Du) \geq \nu|Du|^{p-2}$, for some constants $p \geq 2$ and $\nu > 0$. Let us mention that polyconvex structure (1.17) enjoys (1.26), provided $p \geq 2$, see Sect. 3. Finally, systems in (1.23) and (1.24) satisfy (1.26): details are in Sect. 3.

Let us observe that the interesting Bjorn’s technique allows to deal only with the sub-quadratic case $1 < p \leq 2$. When A_i^α does not depend on u , in Theorem 1.1, we are able to deal with the case $p_0 < p \leq n$, for a suitable $p_0 = p_0(n)$; in the three dimensional case $n = 3$, $p_0 = 3/2$, so our result complements the one of Bjorn and we get boundedness of solutions of elliptic systems under componentwise coercivity, see details at the end of this introduction.

It is worth pointing out that we study system satisfying p, q -growth, according to Marcellini [35]. Regularity in this case is obtained when q is not far from p , see the survey [40] and, more recently, [36, 37, 42]; inequality $p \leq q < p^* \frac{n}{p(n+1)}$ tells us that q cannot be too far from p .

We underline that the strategy for proving our vectorial regularity result is De Giorgi’s elegant and powerful method, see [10]. Precisely, we prove separately that each component u^α satisfies a suitable Caccioppoli-type inequality, a decay of the “excess” on super-(sub-) level sets of u^α that allow to apply iteration arguments and, eventually, the local boundedness of the α -th component of u . A similar strategy has been successfully applied in [6] to prove the boundedness of local minimizers of polyconvex functionals satisfying a non-standard growth, see also [3, 4, 47]. Local boundedness of weak solutions to some elliptic systems with anisotropic or p, q growth has been proved in [7] by using Moser’s iteration technique. In [31, 48], a kind of maximum principle has been proved for systems verifying a condition similar to (1.26); see also [46]. Recent results on the regularity of minimizers of variational integrals or equations in the scalar framework are in [1, 21]; see also [22], where the boundedness of scalar local minimizers of variational integrals is proved under a sharp bound on the exponents p, q , in the light of the counterexamples in [16, 33, 34]. We also cite the interesting paper

[9], where both the scalar and the vectorial case are considered, this last case under the so called Uhlenbeck assumption, i.e. the radial structure $f(x, Du) = \tilde{f}(x, |Du|)$, where f is the energy density of the variational integral. Under this assumption, the usual condition on the exponents p and q to have the local boundedness of vectorial minimizers of functionals, or of weak solutions to systems, is $q < p^*$, see e.g. [7, 8]. Our gap condition on p and q is more restrictive, but we do not require the strong Uhlenbeck assumption.

We try to explain why we are able to consider values of p larger than the ones considered in [2]. Bjorn uses Caccioppoli inequality on superlevel sets $\{v > k\}$ with the *same* exponent p both for Dv and $v - k$. We use Caccioppoli inequality on superlevel sets with *different* exponents: p for Dv and p^* for $v - k$. When p is close to n , then p^* is, by far, larger than p , and this helps a lot. Let us also mention that Bjorn takes $v = \max\{|u^1|, \dots, |u^m|\}$, where $u = (u^1, \dots, u^m)$ is the solution of the system; on the contrary, we take $v = u^\alpha$, the component α of u .

Let us discuss inequalities $1 < p \leq n, p \leq q < p^* \frac{n}{p(n+1)}$, as required in (1.5) of our Theorem 1.1. If $p = n$ the condition on q is trivially satisfied. We have to solve $p < p^* \frac{n}{p(n+1)}$ when $1 < p < n$. This means that $0 < (n + 1)p^2 - n(n + 1)p + n^2$; when $n = 2$ this is satisfied for every p ; when $n = 3$, it is true for $p \neq \frac{3}{2}$; when $n \geq 4$ the inequality is satisfied for $1 < p < p_-$ or $p_+ < p < n$, where

$$p_{\pm} = \frac{n}{2} \left(1 \pm \sqrt{\frac{n-3}{n+1}} \right). \tag{1.27}$$

Note that

$$1 < p_- < 2 < p_+ < n. \tag{1.28}$$

If we confine ourselves to the case $p = q$, it is possible to make a comparison with Bjorn [2]. When $n = 2$, we recover Bjorn’s boundedness result for every $1 < p \leq n = 2$. When $n = 3$, Bjorn’s result is limited to $1 < p \leq 2$ and we complement it, since we are able to deal with $2 < p \leq n = 3$. When $n \geq 4$, Bjorn’s result holds true for $1 < p \leq 2$, our result is valid when $p_+ < p \leq n$, so it remains open the case $2 < p \leq p_+$.

We conclude by observing that in the definition of weak solution of an elliptic equation or system with p, q -growth, the solution is assumed to be in $W_{loc}^{1,q}$ and not in $W_{loc}^{1,p}$, see e.g. [35] and [37]. Enforcing the assumptions on the structure of the nonlinear operator, it is possible to prove the existence of a solution in $W_{loc}^{1,q} \cap (W_0^{1,p} + u_0)$ of a Dirichlet problem with a sufficiently regular boundary datum u_0 , see [5]. On this topic we also refer to Theorem 4.1 in [35], where the scalar case is considered.

Our paper is organized as follows. In the next section we present the proof of Theorem 1.1. In Sect. 3 we give details for some of the previous examples.

2 Proof of Theorem 1.1

The proof of Theorem 1.1 is based on the DeGiorgi method, see [10], suitable for dealing with equations. Nevertheless we apply it in the vectorial framework, since we can apply it to each component u^α of a weak solution u separately.

In what follows we limit ourselves to consider the case $p < n$. The remaining case, $p = n$, can be obtained by the previous one. Indeed, by using the inequality $|z|^{n-\epsilon} - 1 \leq |z|^n$, that holds true for any positive $\epsilon \leq n$, we get that the n, q -growth implies a $n - \epsilon, q$ -growth and

that the assumptions on the exponents, see (1.5), are easily satisfied by choosing $0 < \epsilon < \epsilon_0$, with $\epsilon_0 = \min \left\{ \frac{n^2}{q(n+1)}, n \left(1 - \frac{1}{\tau_1} \right) \right\}$.

STEP 1. Caccioppoli inequality

The particular growth conditions (1.2) and (1.3) guarantee a Caccioppoli inequality for any component u^α of u on every superlevel set $\{u^\alpha > k\}$.

Proposition 2.1 *Let us consider the system (1.1) and assume that (1.2), (1.3) hold. Let $u \in W_{loc}^{1,q}(\Omega; \mathbb{R}^m)$ be a weak solution of (1.1). Let $B_R(x_0) \Subset \Omega$ with $|B_R(x_0)| \leq 1$; for $k \in \mathbb{R}$, $\alpha = 1, \dots, m$ and $0 < \tau \leq R$, denote*

$$A_{k,\tau}^\alpha := \{x \in B_\tau(x_0) : u^\alpha(x) > k\}.$$

If $q \leq p^*$ then, there exists $c = c(n, p, \nu, M) > 0$ such that, for every s, t with $0 < s < t \leq R$, for every $k \in \mathbb{R}$ and for every $\alpha = 1, \dots, m$ we have

$$\begin{aligned} \int_{A_{k,s}^\alpha} |Du^\alpha|^p dx &\leq c \int_{A_{k,t}^\alpha} \left(\frac{u^\alpha - k}{t - s} \right)^{p^*} dx \\ &\quad + c \left\{ \|Du\|_{L^q(B_R(x_0))}^{(q-1)(p^*)'} + \|a\|_{L^1(B_R(x_0))} \right. \\ &\quad \left. + \|b\|_{L^{\tau_2}(B_R(x_0))}^{(p^*)'} \right\} |A_{k,t}^\alpha|^\vartheta, \end{aligned} \tag{2.1}$$

where

$$\vartheta := \min \left\{ 1 - \frac{(p^*)'}{q'}, 1 - \frac{1}{\tau_1}, 1 - \frac{(p^*)'}{\tau_2} \right\}.$$

We can take $c = \frac{1+M2^{1+p^*}}{\nu}$.

Proof Fix $\alpha \in \{1, \dots, m\}$. Consider a cut-off function $\eta \in C_0^1(B_t(x_0))$ satisfying the following assumptions:

$$0 \leq \eta \leq 1, \quad \eta \equiv 1 \text{ in } B_s(x_0), \quad |D\eta| \leq \frac{2}{t - s}. \tag{2.2}$$

Define the test function $\psi = (\psi^1, \dots, \psi^m) \in W_0^{1,q}(B_t(x_0); \mathbb{R}^m)$, where $\psi^\beta = 0$ if $\beta \neq \alpha$ and $\psi^\alpha = (u^\alpha - k)_+ \eta$, where $\tau_+ = \max\{\tau, 0\}$. Notice that

$$\psi_{x_i}^\alpha = \chi_{\{u^\alpha > k\}} u_{x_i}^\alpha \eta + \eta_{x_i} (u^\alpha - k)_+,$$

where $\chi_E(x) = 1$ if $x \in E$ and $\chi_E(x) = 0$ otherwise; moreover, $f_{x_i} = D_i f = \frac{\partial f}{\partial x_i}$.

We insert such a ψ into (1.4) and we get

$$\sum_{i=1}^n \int_{\{u^\alpha > k\}} A_i^\alpha(x, Du) u_{x_i}^\alpha \eta dx = - \sum_{i=1}^n \int_{\{u^\alpha > k\}} A_i^\alpha(x, Du) (u^\alpha - k) \eta_{x_i} dx. \tag{2.3}$$

By (1.2) and (1.3)

$$\begin{aligned}
 \nu \int_{\{u^\alpha > k\}} |Du^\alpha|^p \eta \, dx &\leq \int_{\{u^\alpha > k, \eta > 0\}} a(x) \eta \, dx \\
 &\quad + M \int_{\{u^\alpha > k\}} (u^\alpha - k) |Du|^{q-1} |D\eta| \, dx \\
 &\quad + M \int_{\{u^\alpha > k\}} (u^\alpha - k) b(x) |D\eta| \, dx \\
 &=: J_1 + J_2 + J_3.
 \end{aligned} \tag{2.4}$$

It is easy to estimate J_1 , indeed, using Hölder inequality

$$J_1 \leq \|a\|_{L^1(B_R(x_0))} |A_{k,t}^\alpha|^{1-\frac{1}{\tau_1}}. \tag{2.5}$$

In order to estimate J_2 , we first use Young inequality with exponents p^* and $(p^*)'$.

$$J_2 \leq M \int_{A_{k,t}^\alpha} (u^\alpha - k)^{p^*} |D\eta|^{p^*} \, dx + M \int_{A_{k,t}^\alpha} |Du|^{(q-1)(p^*)'} \, dx.$$

Since $q < p^*$ then $(q - 1)(p^*)' < q$. Therefore we can use Hölder inequality with first exponent $\frac{q'}{(p^*)'} > 1$ to estimate the last integral, obtaining

$$M \int_{A_{k,t}^\alpha} |Du|^{(q-1)(p^*)'} \, dx \leq M \left(\int_{A_{k,t}^\alpha} |Du|^q \, dx \right)^{\frac{(p^*)'}{q'}} |A_{k,t}^\alpha|^{1-\frac{(p^*)'}{q'}}. \tag{2.6}$$

Thus, if we keep in mind that $|D\eta| \leq 2/(t - s)$, then

$$J_2 \leq M 2^{p^*} \int_{A_{k,t}^\alpha} \left(\frac{u^\alpha - k}{t - s} \right)^{p^*} \, dx + M \left(\int_{A_{k,t}^\alpha} |Du|^q \, dx \right)^{\frac{(p^*)'}{q'}} |A_{k,t}^\alpha|^{1-\frac{(p^*)'}{q'}}. \tag{2.7}$$

In order to estimate J_3 , we first use Young inequality with exponents p^* and $(p^*)'$:

$$M \int_{\{u^\alpha > k\}} (u^\alpha - k) b(x) |D\eta| \, dx \leq M \int_{A_{k,t}^\alpha} (u^\alpha - k)^{p^*} |D\eta|^{p^*} \, dx + M \int_{A_{k,t}^\alpha} b^{(p^*)'} \, dx;$$

note that $\tau_2 \geq q' > (p^*)'$; so, we can use Hölder inequality with first exponent $\frac{\tau_2}{(p^*)'} > 1$ and we get

$$M \int_{A_{k,t}^\alpha} b^{(p^*)'} \, dx \leq M \left(\int_{A_{k,t}^\alpha} b(x)^{\tau_2} \, dx \right)^{\frac{(p^*)'}{\tau_2}} |A_{k,t}^\alpha|^{1-\frac{(p^*)'}{\tau_2}}.$$

Once again we use that $|D\eta| \leq 2/(t - s)$, then

$$J_3 \leq M 2^{p^*} \int_{A_{k,t}^\alpha} \left(\frac{u^\alpha - k}{t - s} \right)^{p^*} \, dx + M \left(\int_{B_R(x_0)} b(x)^{\tau_2} \, dx \right)^{\frac{(p^*)'}{\tau_2}} |A_{k,t}^\alpha|^{1-\frac{(p^*)'}{\tau_2}}. \tag{2.8}$$

Collecting (2.4), (2.5), (2.7), (2.8), we get

$$\begin{aligned}
 v \int_{A_{k,t}^\alpha} |Du^\alpha|^p \eta \, dx &\leq M2^{1+p^*} \int_{A_{k,t}^\alpha} \left(\frac{u^\alpha - k}{t - s} \right)^{p^*} dx + \|a\|_{L^{\tau_1}(B_R(x_0))} |A_{k,t}^\alpha|^{1-\frac{1}{\tau_1}} \\
 &\quad + M \left(\int_{A_{k,t}^\alpha} |Du|^q \, dx \right)^{\frac{(p^*)'}{q'}} |A_{k,t}^\alpha|^{1-\frac{(p^*)'}{q'}} \\
 &\quad + M \left(\int_{B_R(x_0)} b(x)^{\tau_2} \, dx \right)^{\frac{(p^*)'}{\tau_2}} |A_{k,t}^\alpha|^{1-\frac{(p^*)'}{\tau_2}} \\
 &\leq M2^{1+p^*} \int_{A_{k,t}^\alpha} \left(\frac{u^\alpha - k}{t - s} \right)^{p^*} dx + \|a\|_{L^{\tau_1}(B_R(x_0))} |A_{k,t}^\alpha|^{1-\frac{1}{\tau_1}} \\
 &\quad + M \|Du\|_{L^q(B_R(x_0))}^{(q-1)(p^*)'} |A_{k,t}^\alpha|^{1-\frac{(p^*)'}{q'}} + M \|b\|_{L^{\tau_2}(B_R(x_0))}^{(p^*)'} |A_{k,t}^\alpha|^{1-\frac{(p^*)'}{\tau_2}}.
 \end{aligned} \tag{2.9}$$

We keep in mind that $\eta = 1$ on $B_s(x_0)$ and $|A_{k,t}^\alpha| \leq |B_R(x_0)| \leq 1$: inequality (2.1) follows by taking $c = \frac{1+M2^{1+p^*}}{v}$. □

STEP 2: Decay of the “excess” on superlevel sets

In this step we consider a scalar Sobolev function $v : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}, n \geq 2$.

Let us assume that Ω is an open set in \mathbb{R}^n and v is a scalar function $v \in W_{loc}^{1,p}(\Omega; \mathbb{R})$, $p \geq 1$. Fix $B_{R_0}(x_0) \Subset \Omega$, with $R_0 < 1$ small enough so that

$$|B_{R_0}(x_0)| < 1 \quad \text{and} \quad \int_{B_{R_0}} |v|^{p^*} \, dx < 1. \tag{2.10}$$

Here $p^* = \frac{np}{n-p}$, since $p < n$.

For every $R \in (0, R_0]$ we define the decreasing sequences

$$\rho_h := \frac{R}{2} + \frac{R}{2^{h+1}} = \frac{R}{2} \left(1 + \frac{1}{2^h} \right), \quad \bar{\rho}_h := \frac{\rho_h + \rho_{h+1}}{2} = \frac{R}{2} \left(1 + \frac{3}{4 \cdot 2^h} \right).$$

Fixed a positive constant $d \geq 1$, define the increasing sequence of positive real numbers

$$k_h := d \left(1 - \frac{1}{2^{h+1}} \right), \quad h \in \mathbb{N}.$$

Moreover, define the sequence $(J_{v,h})$,

$$J_{v,h} := \int_{A_{k_h, \rho_h}} (v - k_h)^{p^*} \, dx,$$

where $A_{k,\rho} = \{v > k\} \cap B_\rho$. The following result holds (see [6, Proposition 2.4], [15, 44]).

Proposition 2.2 *Let $v \in W_{loc}^{1,p}(\Omega; \mathbb{R})$, $p \geq 1$. Fix $B_{R_0}(x_0) \Subset \Omega$, with $R_0 < 1$ small enough such that (2.10) holds. If there exists $0 \leq \vartheta \leq 1$ and $c_0 > 0$ such that for every $0 < s < t \leq R_0$ and for every $k \in \mathbb{R}$*

$$\int_{A_{k,s}} |Dv|^p \, dx \leq c_0 \left\{ \int_{A_{k,t}} \left(\frac{v - k}{t - s} \right)^{p^*} dx + |A_{k,t}|^\vartheta \right\}, \tag{2.11}$$

then, for every $R \in (0, R_0]$,

$$J_{v,h+1} \leq c(\vartheta, R) \left(2 \frac{\rho_h^{p^*}}{\rho_h^p}\right)^h J_{v,h}^{\vartheta},$$

with the positive constant c independent of h .

STEP 3: Iteration and proof of Theorem 1.1

We now resume the proof of Theorem 1.1.

We need the following classical result, see e.g. [17].

Lemma 2.3 *Let $\gamma > 0$ and let (J_h) be a sequence of real positive numbers, such that*

$$J_{h+1} \leq A \lambda^h J_h^{1+\gamma} \quad \forall h \in \mathbb{N} \cup \{0\}, \tag{2.12}$$

with $A > 0$ and $\lambda > 1$. If $J_0 \leq A^{-\frac{1}{\gamma}} \lambda^{-\frac{1}{\gamma^2}}$, then $J_h \leq \lambda^{-\frac{h}{\gamma}} J_0$ and $\lim_{h \rightarrow \infty} J_h = 0$.

Fix $B_{R_0}(x_0) \Subset \Omega$, with $R_0 < 1$ small enough such that $|B_{R_0}(x_0)| < 1$ and $\int_{B_{R_0}} |u|^{p^*} dx < 1$. By Proposition 2.1 we have that u^α satisfies (2.1); i.e. for every $0 < s < t \leq R_0$ and every $k \in \mathbb{R}$,

$$\begin{aligned} \int_{A_{k,s}^\alpha} |Du^\alpha|^p dx &\leq c \int_{A_{k,t}^\alpha} \left(\frac{u^\alpha - k}{t - s}\right)^{p^*} dx \\ &+ c \left\{ \|Du\|_{L^q(B_{R_0}(x_0))}^{(q-1)(p^*)'} + \|a\|_{L^{\tau_1}(B_{R_0}(x_0))} \right. \\ &\left. + \|b\|_{L^{\tau_2}(B_{R_0}(x_0))}^{(p^*)'} \right\} |A_{k,t}^\alpha|^\vartheta, \end{aligned}$$

where

$$\vartheta := \min \left\{ 1 - \frac{(p^*)'}{q'}, 1 - \frac{1}{\tau_1}, 1 - \frac{(p^*)'}{\tau_2} \right\}$$

and $c = \frac{1+M2^{1+p^*}}{\nu}$.

Therefore the scalar function u^α satisfies (2.11) of Proposition 2.2 with constant c_0 depending on

$$\|Du\|_{L^q(B_{R_0}(x_0))}^{(q-1)(p^*)'}, \quad \|a\|_{L^{\tau_1}(B_{R_0}(x_0))} \quad \text{and} \quad \|b\|_{L^{\tau_2}(B_{R_0}(x_0))}^{(p^*)'}.$$

Note that these integrals are finite.

As above, let us define

$$k_h := d \left(1 - \frac{1}{2^{h+1}}\right), \quad h \in \mathbb{N}$$

with $d \geq 1$ (d will be fixed later) and, for every $R \in (0, R_0]$, define

$$\rho_h := \frac{R}{2} + \frac{R}{2^{h+1}} = \frac{R}{2} \left(1 + \frac{1}{2^h}\right), \quad \bar{\rho}_h := \frac{\rho_h + \rho_{h+1}}{2} = \frac{R}{2} \left(1 + \frac{3}{4 \cdot 2^h}\right)$$

and

$$J_{u^\alpha,h} := \int_{A_{k_h,\rho_h}^\alpha} (u^\alpha - k_h)^{p^*} dx.$$

Proposition 2.2, applied to u^α , gives

$$J_{u^\alpha, h+1} \leq c(\vartheta, R) \left(2^{\frac{p^* p^*}{p}}\right)^h J_{u^\alpha, h}^{\frac{p^*}{p}}, \tag{2.13}$$

with the positive constant c independent of h and, by (1.5), with the exponent $\vartheta \frac{p^*}{p}$ greater than 1. Indeed, we notice that $q < p^* \frac{n}{p(n+1)}$ is equivalent to $\frac{q}{q-1} > \frac{p^*}{p^*-1} \frac{n}{p}$; therefore (1.5) implies

$$\frac{p}{p^*} < \min \left\{ 1 - \frac{(p^*)'}{q'}, 1 - \frac{1}{\tau_1}, 1 - \frac{(p^*)'}{\tau_2} \right\} = \vartheta,$$

so we get $1 < \vartheta \frac{p^*}{p}$.

Moreover, since

$$J_{u^\alpha, 0} = \int_{A_{\frac{d}{2}, R}^\alpha} \left(u^\alpha - \frac{d}{2}\right)^{p^*} dx \rightarrow 0 \text{ as } d \rightarrow +\infty,$$

we can choose $d \geq 1$ large enough, so that

$$J_{u^\alpha, 0} < c(\vartheta, R)^{-\frac{1}{\vartheta \frac{p^*}{p} - 1}} \left(2^{\frac{p^* p^*}{p}}\right)^{-\frac{1}{(\vartheta \frac{p^*}{p} - 1)^2}}.$$

Therefore, by Lemma 2.3, $\lim_{h \rightarrow +\infty} J_{u^\alpha, h} = 0$. Thus, $u^\alpha \leq d$ a.e. in $B_{\frac{R}{2}}(x_0)$. We have so proved that u^α is locally bounded from above.

To prove that u^α is locally bounded from below, we can observe that $\tilde{u} = -u$ is a weak solution for

$$\sum_{i=1}^n \frac{\partial}{\partial x_i} \left(\tilde{A}_i^\alpha(x, D\tilde{u}(x))\right) = 0, \quad 1 \leq \alpha \leq m,$$

where $\tilde{A}(x, z) = -A(x, -z)$. It is easy to check that \tilde{A} satisfies assumptions analogous to (1.2) and (1.3). Therefore, by what previously proved, there exists d' such that $\tilde{u}^\alpha = -u^\alpha \leq d'$ a.e. in $B_{\frac{R}{2}}(x_0)$. We have so proved that $u^\alpha \in L^\infty(B_{\frac{R}{2}}(x_0))$. Due to the arbitrariness of x_0 and R_0 , we get $u^\alpha \in L_{loc}^\infty(\Omega)$.

3 Examples

Example 1 We consider example (1.17) that we rewrite for the convenience of the reader:

$$A_i^\alpha(z) = p|z|^{p-2} z_i^\alpha + h'(\det z)(\text{Cof } z)_i^\alpha, \tag{3.1}$$

where $m = n$, $z \in \mathbb{R}^{n \times n}$, $\det z = \sum_{i=1}^n z_i^\alpha (\text{Cof } z)_i^\alpha$; moreover, h is convex, bounded from below and C^1 . Exploiting the convexity of h , we get

$$h(0) \geq h(t) + h'(t)(0 - t), \tag{3.2}$$

so that

$$h'(t)t \geq h(t) - h(0) \geq \inf_{\mathbb{R}} h - h(0). \tag{3.3}$$

Let us compute the indicator function for this choice of A : we get

$$\begin{aligned}
 I_A(x, u, z) &= \sum_{i,\alpha,\beta} A_i^\alpha(z) z_i^\beta \frac{u^\alpha u^\beta}{|u|^2} = \sum_{i,\alpha,\beta} p|z|^{p-2} z_i^\alpha z_i^\beta \frac{u^\alpha u^\beta}{|u|^2} \\
 &\quad + \sum_{i,\alpha,\beta} h'(\det z) (\text{Cof } z)_i^{\alpha\beta} z_i^\alpha z_i^\beta \frac{u^\alpha u^\beta}{|u|^2} \\
 &= p|z|^{p-2} \sum_i \sum_\alpha z_i^\alpha \frac{u^\alpha}{|u|} \sum_\beta z_i^\beta \frac{u^\beta}{|u|} + h'(\det z) \sum_{\alpha,\beta} \frac{u^\alpha u^\beta}{|u|^2} \sum_i (\text{Cof } z)_i^{\alpha\beta} z_i^\alpha z_i^\beta \\
 &= p|z|^{p-2} \sum_i \left(\sum_\alpha z_i^\alpha \frac{u^\alpha}{|u|} \right)^2 + h'(\det z) \sum_\alpha \frac{u^\alpha u^\alpha}{|u|^2} \sum_i (\text{Cof } z)_i^{\alpha\alpha} z_i^\alpha z_i^\alpha \\
 &= p|z|^{p-2} \sum_i \left(\sum_\alpha z_i^\alpha \frac{u^\alpha}{|u|} \right)^2 + h'(\det z) \det z \geq \inf_{\mathbb{R}} h - h(0),
 \end{aligned}$$

where we used the property $\sum_i (\text{Cof } z)_i^{\alpha\beta} z_i^\alpha z_i^\beta = 0$ if $\beta \neq \alpha$. When $h(0) = \inf_{\mathbb{R}} h$, then strong Meier’s condition (1.11) is satisfied; if $h(0) > \inf_{\mathbb{R}} h$, then weak Meier’s condition (1.19) is verified with $\lambda = 1$, $d(x) = 0$ and $g(x) = h(0) - \inf_{\mathbb{R}} h$. Now, let us verify componentwise coercivity (1.26). We have

$$\begin{aligned}
 \sum_i A_i^\alpha(z) z_i^\alpha &= \sum_i p|z|^{p-2} z_i^\alpha z_i^\alpha + \sum_i h'(\det z) (\text{Cof } z)_i^{\alpha\alpha} z_i^\alpha z_i^\alpha \\
 &= p|z|^{p-2} |z^\alpha|^2 + h'(\det z) \det z \geq p|z^\alpha|^p + \inf_{\mathbb{R}} h - h(0),
 \end{aligned}$$

provided $p \geq 2$; then (1.26) is verified with $\nu = p$, $a(x) = h(0) - \inf_{\mathbb{R}} h$ and $b(x) = 0$.

Example 2 We consider example (1.20) that we rewrite for the convenience of the reader:

$$A_i^\alpha(x, z) = \sigma^\alpha(x) z_i^\alpha, \tag{3.4}$$

where $m = 2$, $\sigma^1(x) = 18 + 2 \sin(|x|^2)$ and $\sigma^2(x) = 2 + \sin(|x|^2)$. Since $\sigma^\alpha(x) \geq 1$, it is easy to check (1.26):

$$\sum_i A_i^\alpha(x, z) z_i^\alpha = \sum_i \sigma^\alpha(x) z_i^\alpha z_i^\alpha = \sigma^\alpha(x) |z^\alpha|^2 \geq |z^\alpha|^2;$$

so, (1.26) is verified with $p = 2$, $\nu = 1$, $a(x) = 0$ and $b(x) = 0$. We are going to show that (1.19) is not fulfilled. Indeed, we take $u^1 = u^2 = s > 0$ with s large enough (see (3.5) later); moreover, we take $z_i^\alpha = 0$ if $i \geq 2$, $z_1^1 = -s^2$, $z_1^2 = 2s^2$. Then $|z|^2 = 5s^4$, $|u|^2 = 2s^2$, $\frac{u^\alpha u^\beta}{|u|^2} = \frac{1}{2}$ and

$$\begin{aligned}
 \sum_{i,\alpha,\beta} A_i^\alpha(x, z) z_i^\beta \frac{u^\alpha u^\beta}{|u|^2} &= \frac{1}{2} \sum_{\alpha,\beta} A_1^\alpha(x, z) z_1^\beta = \frac{1}{2} \sum_\alpha A_1^\alpha(x, z) \sum_\beta z_1^\beta \\
 &= \frac{1}{2} (\sigma^1(x) z_1^1 + \sigma^2(x) z_1^2) (z_1^1 + z_1^2) = \frac{1}{2} (-\sigma^1(x) + 2\sigma^2(x)) s^4 = -7s^4 \underbrace{\leq}_{0 < \delta < 1} -\delta 7s^4
 \end{aligned}$$

$$\begin{aligned}
 &= -\delta s^4 \{5 + 1 + 1\} \underbrace{\leq}_{(*)} -\delta s^4 \left\{ 5 + \left(\frac{1}{\delta}\right)^{\lambda+1} \frac{2d(x)}{s^2} + \left(\frac{1}{\delta}\right)^{\lambda+1} \frac{g(x)}{s^4} \right\} \\
 &= - \left\{ \delta |z|^2 + \left(\frac{1}{\delta}\right)^\lambda [d(x)|u|^2 + g(x)] \right\},
 \end{aligned}$$

where (*) is guaranteed by the choice of s as follows

$$s = \max \left\{ L; \left[\left(\frac{1}{\delta}\right)^{\lambda+1} 2d(x) \right]^{1/2}; \left[\left(\frac{1}{\delta}\right)^{\lambda+1} g(x) \right]^{1/4} \right\}. \tag{3.5}$$

Example 3 Let us consider example (1.23) that we rewrite for the convenience of the reader:

$$A_i^\alpha(z) = 2z_i^\alpha + h'(z_1^\alpha z_1^2) z_1^{\hat{\alpha}} \delta_{i1}, \tag{3.6}$$

where $m = 2, \hat{\alpha} = 2$ if $\alpha = 1$ and $\hat{\alpha} = 1$ if $\alpha = 2$; moreover, $\delta_{i1} = 1$ if $i = 1$ and $\delta_{i1} = 0$ otherwise. Here, h is convex, C^1 , bounded from below and $h'(0) \leq -8$. For instance,

$$h(t) = 16\sqrt{1 + (t - 1)^2}. \tag{3.7}$$

Let us first check (1.26):

$$\begin{aligned}
 \sum_i A_i^\alpha(z) z_i^\alpha &= \sum_i 2z_i^\alpha z_i^\alpha + \sum_i h'(z_1^\alpha z_1^2) z_1^{\hat{\alpha}} \delta_{i1} z_i^\alpha \\
 &= 2|z^\alpha|^2 + h'(z_1^\alpha z_1^2) z_1^{\hat{\alpha}} z_1^\alpha = 2|z^\alpha|^2 + h'(z_1^\alpha z_1^2) z_1 z_1^2 \geq 2|z^\alpha|^2 + \inf_{\mathbb{R}} h - h(0),
 \end{aligned}$$

since $z_1^{\hat{\alpha}} z_1^\alpha = z_1 z_1^2$; then (1.26) is verified with $v = 2, p = 2, a(x) = h(0) - \inf_{\mathbb{R}} h$ and $b(x) = 0$. We are going to show that (1.19) is not fulfilled. Indeed, we take $u^1 = u^2 = s > 0$ with s large enough (see (3.5) as before); moreover, we take $z_1^2 = s^2$ and $z_i^\alpha = 0$ otherwise. Then $|z|^2 = s^4, |u|^2 = 2s^2, \frac{u^\alpha u^\beta}{|u|^2} = \frac{1}{2}$ and

$$\begin{aligned}
 \sum_{i,\alpha,\beta} A_i^\alpha(z) z_i^\beta \frac{u^\alpha u^\beta}{|u|^2} &= \frac{1}{2} \sum_{\alpha,\beta} A_1^\alpha(z) z_1^\beta = \frac{1}{2} \sum_{\alpha} A_1^\alpha(z) \sum_{\beta} z_1^\beta \\
 &= \frac{1}{2} (2z_1^2 + h'(0)z_1^2)(z_1^2) = \frac{1}{2} (2 + h'(0))s^4 \underbrace{\leq}_{h'(0) \leq -8} \frac{1}{2} (2 - 8)s^4 = -3s^4 \underbrace{\leq}_{0 < \delta < 1} -\delta 3s^4 \\
 &= -\delta s^4 \{1 + 1 + 1\} \underbrace{\leq}_{(*)} -\delta s^4 \left\{ 1 + \left(\frac{1}{\delta}\right)^{\lambda+1} \frac{2d(x)}{s^2} + \left(\frac{1}{\delta}\right)^{\lambda+1} \frac{g(x)}{s^4} \right\} \\
 &= - \left\{ \delta |z|^2 + \left(\frac{1}{\delta}\right)^\lambda [d(x)|u|^2 + g(x)] \right\},
 \end{aligned}$$

where (*) is guaranteed by the choice of s (3.5) as before. In order to show that we can use Theorem 1.1, we use formula (3.7) and we select $n = 3$. Then $|h'(t)| \leq 16$ and we get

$$\sum_{i=1}^3 |A_i^\alpha(z)| \leq 54|z|, \tag{3.8}$$

so (1.3) is satisfied with $q = 2, M = 54$ and $b(x) = 0, \tau_2 = +\infty$. Note that previous calculations checked the validity of (1.2) with $p = 2, v = 2, a(x) = h(0) - \inf_{\mathbb{R}} h = 16(\sqrt{2} - 1)$ and $\tau_1 = +\infty$. Since we selected $n = 3, q = p = 2$, then $\frac{3}{2} = p_0(n) < 2 = q = p < p^* = \frac{n}{p(n+1)}$; this implies that (1.5) is satisfied and we can use our Theorem 1.1 and we get the following

Corollary 3.1 *If Ω is a bounded open subset of \mathbb{R}^3 , then all solutions $u \in W_{loc}^{1,2}(\Omega; \mathbb{R}^2)$ of system (1.1), with $n = 3, m = 2$, (3.6) and (3.7), are locally bounded in Ω .*

Example 4 Let us consider example (1.24) that we rewrite for the convenience of the reader:

$$A_i^\alpha(x, z) = \sigma^\alpha(x)p|z|^{p-2}z_i^\alpha + h'(z_1^1 z_1^2)z_1^{\hat{\alpha}}\delta_{i1}, \tag{3.9}$$

where $m = 2, \sigma^1(x) = 48 + 3 \sin(|x|^2)$ and $\sigma^2(x) = 2 + \sin(|x|^2), \hat{\alpha} = 2$ if $\alpha = 1$ and $\hat{\alpha} = 1$ if $\alpha = 2$; moreover, $\delta_{i1} = 1$ if $i = 1$ and $\delta_{i1} = 0$ otherwise. Here, $2 \leq p, h$ is convex, C^1 , bounded from below and $h'(0) \leq 0$. For instance,

$$h(t) = (1 + t^2)^{p/4}. \tag{3.10}$$

Let us first check (1.26); since $\sigma^\alpha(x) \geq 1$ and $z_1^{\hat{\alpha}}z_1^\alpha = z_1^1 z_1^2$,

$$\begin{aligned} \sum_i A_i^\alpha(x, z)z_i^\alpha &= \sum_i \sigma^\alpha(x)p|z|^{p-2}z_i^\alpha z_i^\alpha + \sum_i h'(z_1^1 z_1^2)z_1^{\hat{\alpha}}\delta_{i1}z_i^\alpha \\ &= \sigma^\alpha(x)p|z|^{p-2}|z^\alpha|^2 + h'(z_1^1 z_1^2)z_1^{\hat{\alpha}}z_1^\alpha = \sigma^\alpha(x)p|z|^{p-2}|z^\alpha|^2 + h'(z_1^1 z_1^2)z_1^1 z_1^2 \\ &\geq p|z^\alpha|^p + \inf_{\mathbb{R}} h - h(0), \end{aligned}$$

then (1.26) is verified with $v = p, a(x) = h(0) - \inf_{\mathbb{R}} h$ and $b(x) = 0$. We are going to show that (1.19) is not fulfilled. Indeed, we take $u^1 = u^2 = s > 0$ with s large enough (see (3.11) later); moreover, we take $z_1^1 = s^2, z_1^2 = 0, z_2^1 = -2s^2, z_2^2 = 3s^2$ and $z_i^\alpha = 0$ otherwise. Then $|z|^2 = 14s^4, |u|^2 = 2s^2, \frac{u^\alpha u^\beta}{|u|^2} = \frac{1}{2}$ and

$$\begin{aligned} \sum_{i,\alpha,\beta} A_i^\alpha(x, z)z_i^\beta \frac{u^\alpha u^\beta}{|u|^2} &= \frac{1}{2} \sum_{i,\alpha,\beta} A_i^\alpha(x, z)z_i^\beta = \frac{1}{2} \sum_i \sum_\alpha A_i^\alpha(x, z) \sum_\beta z_i^\beta \\ &= \frac{1}{2} \sum_i \sum_\alpha \sigma^\alpha(x)p|z|^{p-2}z_i^\alpha \sum_\beta z_i^\beta + \frac{1}{2} \sum_i \sum_\alpha h'(z_1^1 z_1^2)z_1^{\hat{\alpha}}\delta_{i1} \sum_\beta z_i^\beta \\ &= \frac{1}{2} \sum_i \sum_\alpha \sigma^\alpha(x)p|z|^{p-2}z_i^\alpha \sum_\beta z_i^\beta + \frac{1}{2} \sum_\alpha h'(z_1^1 z_1^2)z_1^{\hat{\alpha}} \sum_\beta z_1^\beta \\ &= \frac{p}{2} |z|^{p-2} \sum_i \{[\sigma^1(x)z_i^1 + \sigma^2(x)z_i^2][z_i^1 + z_i^2] + \frac{1}{2}h'(z_1^1 z_1^2)[z_1^2 + z_1^1][z_1^1 + z_1^2]\} \\ &= \frac{p}{2} |z|^{p-2} \{[\sigma^1(x)z_1^1 + \sigma^2(x)z_1^2][z_1^1 + z_1^2] + [\sigma^1(x)z_2^1 + \sigma^2(x)z_2^2][z_2^1 + z_2^2]\} \\ &\quad + \frac{1}{2}h'(z_1^1 z_1^2)[z_1^1 + z_1^2]^2 \\ &= \frac{p}{2} |z|^{p-2} \{ \sigma^1(x)s^4 + [-2\sigma^1(x) + 3\sigma^2(x)]s^4 + \frac{1}{2}h'(0)s^4 \} \\ &\stackrel{h'(0) \leq 0}{\leq} \frac{p}{2} |z|^{p-2} s^4 \{-\sigma^1(x) + 3\sigma^2(x)\} = \frac{p}{28} |z|^p \{-\sigma^1(x) + 3\sigma^2(x)\} = -3\frac{p}{2} |z|^p \end{aligned}$$

$$\begin{aligned} & \underbrace{\leq}_{2 \leq p} -3|z|^p \underbrace{\leq}_{0 < \delta < 1} -\delta|z|^p 3 = -\delta|z|^p \{1 + 1 + 1\} \\ & \underbrace{\leq}_{(**)} -\delta|z|^p \left\{ 1 + \left(\frac{1}{\delta}\right)^{\lambda+1} d(x) \frac{1}{(\sqrt{7}s)^p} + \left(\frac{1}{\delta}\right)^{\lambda+1} \frac{g(x)}{(\sqrt{14}s^2)^p} \right\} \\ & = - \left\{ \delta|z|^p + \left(\frac{1}{\delta}\right)^\lambda [d(x)|u|^p + g(x)] \right\}, \end{aligned}$$

where (**) is guaranteed by the choice (3.11) of s as follows.

$$s = \max \left\{ L; \left[\left(\frac{1}{\delta}\right)^{\lambda+1} \frac{d(x)}{(\sqrt{7})^p} \right]^{1/p}; \left[\left(\frac{1}{\delta}\right)^{\lambda+1} \frac{g(x)}{(\sqrt{14})^p} \right]^{1/(2p)} \right\}. \tag{3.11}$$

In order to show that we can use Theorem 1.1, we use formula (3.10), we select $n = 3$ and we require $p < 3 = n$. Then $|h'(t)| \leq \frac{p}{2} (1 + t^2)^{\frac{p-2}{4}}$ and we get

$$\begin{aligned} \sum_{i=1}^3 |A_i^\alpha(x, z)| & \leq 3\sigma^\alpha(x)p|z|^{p-1} + 3\frac{p}{2} (1 + |z|^4)^{\frac{p-2}{4}} |z| \leq 153p|z|^{p-1} \\ & + 3\frac{p}{2} (1 + |z|^4)^{\frac{p-1}{4}} \\ & \leq 153p|z|^{p-1} + 3\frac{p}{2} 2^{\frac{p-1}{4}} (1 + |z|^{p-1}) \leq p(153 + 2^{1+\frac{p-1}{4}})(|z|^{p-1} + 1), \end{aligned}$$

so (1.3) is satisfied with $q = p$, $M = p(153 + 2^{1+\frac{p-1}{4}})$ and $b(x) = 1$, $\tau_2 = +\infty$. Note that previous calculations checked the validity of (1.2) with $v = p$, $a(x) = h(0) - \inf_{\mathbb{R}} h = 0$ and $\tau_1 = +\infty$. Since we selected $n = 3$, $q = p \in [2, 3)$, then $\frac{3}{2} = p_0(n) < 2 \leq q = p < p^* \frac{n}{p(n+1)}$; this implies that (1.5) is satisfied and we can use our Theorem 1.1 and we get the following

Corollary 3.2 *If Ω is a bounded open subset of \mathbb{R}^3 , then all solutions $u \in W_{loc}^{1,p}(\Omega; \mathbb{R}^2)$ of system (1.1), with $2 \leq p < 3 = n$, $m = 2$, (3.9) and (3.10), are locally bounded in Ω .*

Acknowledgements The authors thank the referees for useful remarks. The authors are members of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). The authors gladly take the opportunity to thank GNAMPA, INdAM, UNIBO, UNIFI, UNIVAQ for the support.

Data availability statement data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Bella, P., Schäffner, M.: On the regularity of minimizers for scalar integral functionals with (p, q) -growth. *Anal. PDE* **13**(7), 2241–2257 (2020)
2. Bjorn, J.: Boundedness and differentiability for nonlinear elliptic systems. *Trans. Am. Math. Soc.* **353**(11), 4545–4565 (2001)
3. Carozza, M., Gao, H., Giova, R., Leonetti, F.: A boundedness result for minimizers of some polyconvex integrals. *J. Opt. Theory Appl.* **178**(3), 699–725 (2018)
4. Cupini, G., Focardi, M., Leonetti, F., Mascolo, E.: On the Hölder continuity for a class of vectorial problems. *Adv. Nonlinear Anal.* **9**(1), 1008–1025 (2020)
5. Cupini, G., Leonetti, F., Mascolo, E.: Existence of weak solutions for elliptic systems with p, q -growth. *Ann. Acad. Sci. Fenn. Math.* **40**(2), 645–658 (2015)
6. Cupini, G., Leonetti, F., Mascolo, E.: Local boundedness for minimizers of some polyconvex integrals. *Arch. Ration. Mech. Anal.* **224**, 269–289 (2017)
7. Cupini, G., Marcellini, P., Mascolo, E.: Local boundedness of solutions to quasilinear elliptic systems. *Manuscripta Math.* **137**, 287–315 (2012)
8. Cupini, G., Marcellini, P., Mascolo, E.: Local boundedness of solutions to some anisotropic elliptic systems, *Recent trends in nonlinear partial differential equations. II. Stationary problems*, 169–186, *Contemp. Math.*, 595, Amer. Math. Soc., Providence, RI, (2013)
9. De Filippis, C., Mingione, G.: Lipschitz bounds and nonautonomous integrals. *Arch. Ration. Mech. Anal.* **242**(2), 973–1057 (2021)
10. De Giorgi, E.: Sulla differenziabilità e l'analiticità delle estremali degli integrali multipli regolari. *Mem. Accad. Sci. Torino. Cl. Sci. Fis. Mat. Nat.* **3**, 25–43 (1957)
11. De Giorgi, E.: Un esempio di estremali discontinue per un problema variazionale di tipo ellittico. *Boll. Un. Mat. Ital.* **4**(1), 135–137 (1968)
12. Fattorusso, L., Softova, L.: Precise Morrey regularity of the weak solutions to a kind of quasilinear systems with discontinuous data, *Electron. J. Qual. Theory Differ. Equ.*, Paper No. 36, 13 pp (2020)
13. Frehse, J.: Una generalizzazione di un controesempio di De Giorgi nella teoria delle equazioni ellittiche. *Boll. Un. Mat. Ital.* (4) **3**, 998–1002 (1970)
14. Frehse, J.: An irregular complex valued solution to a scalar uniformly elliptic equation. *Calc. Var. Partial Differ. Equ.* **33**, 263–266 (2008)
15. Fusco, N., Sbordone, C.: Some remarks on the regularity of minima of anisotropic integrals, *Commu. P. D. E.*, **18**, 153–167 (1993)
16. Giaquinta, M.: Growth conditions and regularity, a counterexample. *Manuscripta Math.* **59**, 245–248 (1987)
17. Giusti, E.: *Direct Methods in the Calculus of Variations*. World Scientific Publishing Co., Inc., River Edge (2003)
18. Giusti, E., Miranda, M.: Un esempio di soluzioni discontinue per un problema di minimo relativo ad un integrale regolare del calcolo delle variazioni. *Boll. Un. Mat. Ital.* **2**, 1–8 (1968)
19. Hao, W., Leonardi, S., Necas, J.: An example of irregular solution to a nonlinear Euler-Lagrange elliptic system with real analytic coefficients. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4) **23**(1), 57–67 (1996)
20. Hao, W., Leonardi, S., Steinhauer, M.: Examples of discontinuous, divergence-free solutions to elliptic variational problems. *Comment. Math. Univ. Carolin.* **36**(3), 511–517 (1995)
21. Hästö, P., Ok, J.: Regularity theory for non-autonomous partial differential equations without Uhlenbeck structure, *Preprint* (2021), [arXiv:2110.14351](https://arxiv.org/abs/2110.14351)
22. Hirsch, J., Schäffner, M.: Growth conditions and regularity, an optimal local boundedness result, *Commun. Contemp. Math.* **23**, no. 3, Paper No. 2050029, 17 pp (2021)
23. John, O., Maly, J., Stara, J.: Nowhere continuous solutions to elliptic systems. *Comment. Math. Univ. Carolin.* **30**(1), 33–43 (1989)
24. John, O., Necas, J., Stara, J.: Counterexample to the regularity of weak solution of elliptic systems. *Comment. Math. Univ. Carolin.* **21**(1), 145–154 (1980)
25. Kristensen, J., Mingione, G.: Sketches of Regularity Theory from The 20th Century and the Work of Jindrich Necas. In: “Selected works of Jindrich Necas”. PDEs, continuum mechanics and regularity. Edited by Sarka Necasova, Milan Pokorny and Vladimir Sverak. *Advances in Mathematical Fluid Mechanics. Birkhauser/Springer, Basel*, (2015). xiv+785 pp
26. Krömer, S.: A priori estimates in L^∞ for non-diagonal perturbed quasilinear systems. *Ann. Sc. Norm. Super. Pisa Cl. Sci.* (5) **8**, 417–428 (2009)
27. Ladyzhenskaya, O., Ural' tseva, N.: *Linear and Quasilinear Elliptic Equations*. Academic Press, New York (1968)

28. Landes, R.: Some remarks on bounded and unbounded weak solutions of elliptic systems. *Manuscripta Math.* **64**, 227–234 (1989)
29. Landes, R.: On the regularity of weak solutions of certain elliptic systems. *Calc. Var. Partial Differ. Equ.* **25**, 247–255 (2005)
30. Leonetti, F., Mascolo, E.: Local boundedness for vector valued minimizers of anisotropic functionals. *Z. Anal. Anwend.* **31**, 357–378 (2012)
31. Leonetti, F., Petricca, P.V.: Regularity for solutions to some nonlinear elliptic systems. *Complex Var. Elliptic Equ.* **56**(12), 1099–1113 (2011)
32. Leonetti, F., Petricca, P.V.: Summability for solutions to some quasilinear elliptic systems. *Ann. Mat. Pura Appl.* **193**, 1671–1682 (2014)
33. Marcellini, P.: Un exemple de solution discontinue d'un problème variationnel dans le cas scalaire, *Preprint 11, Istituto Matematico "U.Dini"*, Università di Firenze, (1987)
34. Marcellini, P.: Regularity of minimizers of integrals of the calculus of variations with nonstandard growth conditions. *Arch. Rational Mech. Anal.* **105**(3), 267–284 (1989)
35. Marcellini, P.: Regularity and existence of solutions of elliptic equations with p, q -growth conditions. *J. Differ. Equ.* **90**(1), 1–30 (1991)
36. Marcellini, P.: Regularity under general and p, q -growth conditions. *Discrete Cont. Dyn. Syst. Ser. S* **13**, 2009–2031 (2020)
37. Marcellini, P.: Growth conditions and regularity for weak solutions to nonlinear elliptic PDES. *J. Math. Anal. Appl.* **501**(1), 124408 (2021)
38. Maz'ja, V.G.: Examples of nonregular solutions of quasilinear elliptic equations with analytic coefficients. (Russian) *Funkcional Anal. i Prilozen* **2**(3), 53–57 (1968)
39. Meier, M.: Boundedness and integrability properties of weak solutions of quasilinear elliptic systems. *J. Reine Angew. Math.* **333**, 191–220 (1982)
40. Mingione, G.: Regularity of minima: an invitation to the dark side of the calculus of variations. *Appl. Math.* **51**, 355–426 (2006)
41. Mingione, G.: Singularities of minima: a walk on the wild side of the calculus of variations. *J. Global Optim.* **40**, 209–223 (2008)
42. Mingione, G., Radulescu, V.: Recent developments in problems with nonstandard growth and nonuniform ellipticity. *J. Math. Anal. Appl.* **501**(1), 125197 (2021)
43. Mooney, C., Savin, O.: Some singular minimizers in low dimensions in the calculus of variations. *Arch. Ration. Mech. Anal.* **221**, 1–22 (2016)
44. Moscarriello, G., Nania, L.: Hölder continuity of minimizers of functionals with non standard growth conditions. *Ricerche Mat.* **40**, 259–273 (1991)
45. Nečas, J.: Example of an irregular solution to a nonlinear elliptic system with analytic coefficients and conditions for regularity, *Theory Nonlin. Oper.*, Abhand. der Wiss. der DDR (1977)
46. Palagachev, D., Softova, L.: Boundedness of solutions to a class of coercive systems with Morrey data. *Nonlinear Anal.* **191**, 111630 (2020)
47. Shan, Y., Gao, H.: Holder continuity for vectorial local minimizers of variational integrals, *Rocky Mountain Math. J.* online-first <https://projecteuclid.org/journals/rmj/rocky-mountain-journal-of-mathematics/DownloadAcceptedPapers/210524-HongyaGao.pdf>
48. Softova, L.: Boundedness of the solutions to non-linear systems with Morrey data. *Complex Var. Elliptic Equ.* **63**(11), 1581–1594 (2018)
49. Soucek, J.: Singular solutions to linear elliptic systems. *Comment. Math. Univ. Carolin.* **25**(2), 273–281 (1984)
50. Sverák, V., Yan, X.: A singular minimizer of a smooth strongly convex functional in three dimensions. *Calc. Var. Partial Differ. Equ.* **10**(3), 213–221 (2000)